

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 413/04, 417/04, 239/54, A01N 43/78, 43/76		A1	(11) International Publication Number: WO 97/08170
			(43) International Publication Date: 6 March 1997 (06.03.97)
(21) International Application Number: PCT/US96/13995		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 30 August 1996 (30.08.96)			
(30) Priority Data: 60/003,080 31 August 1995 (31.08.95) US Not furnished 31 July 1996 (31.07.96) US			
(71) Applicant: FMC CORPORATION [US/US]; 1735 Market Street, Philadelphia, PA 19103 (US).			
(72) Inventors: CRAWFORD, Scott, D.; 130 Birch Hollow Drive, Bordentown, NJ 08505 (US). MARAVETZ, Lester, L.; 843 Carlton Road, Westfield, NJ 07090 (US). THEODORIDIS, George; 45 Monroe Lane, Princeton, NJ 08540 (US).			
(74) Agent: KENNEDY, Robert, M.; FMC Corporation, 1735 Market Street, Philadelphia, PA 19103 (US).		Published With international search report.	

(54) Title: HERBICIDAL 3-(SUBSTITUTED BENZOXAZOL-7-YL) AND 3-(SUBSTITUTED BENZOTHIAZOL-7-YL)-1-SUBSTITUTED-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONES



ALKENYL⁶, ALKYNYL⁶, ARYL⁶, ARYLALKYL⁶, ALKYLARYLALKYL⁶, HALOALKYL⁶,
HYDROXY, ALKOXY, HYDROXYALKYL⁶, HALOARYL⁶, HALOARYLALKYL⁶,
ALKOXYARYL⁶, SULFHYDRYL⁶, ALKYLTHIO, PIPERIDINYL⁶, ALKYLAMINO,
ALKOXYALKYL⁶, PHENOXY, AMINO, ALKYL-SULFONYLAMINO, ARYL-SULFONYLAMINO

(57) Abstract

Herbicidal 3-(substituted benzoxazol-7-yl) and 3-(substituted benzothiazol-7-yl)-1-substituted-6-trifluoromethyl-2,4-(1H,3H)pyrimidinediones, compositions containing them, and methods of using them to control undesired plant growth are disclosed, as are novel intermediates used in the preparation of these compounds. The herbicidal compounds of the present invention are defined by generic structure α , in which R is selected from a variety of substituents, including halogen, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkylarylalkyl, haloalkyl, hydroxy, alkoxy, hydroxyalkyl, haloaryl, haloarylalkyl, alkoxyaryl, sulfhydryl, alkylthio, piperidinyl, alkylamino, alkoxyalkyl, phenoxy, amino, alkylsulfonamino, arylsulfonamino, and carboxy; R¹ is alkyl or amino; R² is hydrogen or halogen; X is oxygen or sulfur; Y is hydrogen, halogen, alkoxy, cyano, or nitro; and Z is halogen; where halogen is bromine, chlorine, fluorine, or iodine, and each alkyl moiety has one to six carbon atoms.

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**HERBICIDAL 3-(SUBSTITUTED BENZOXAZOL-7-YL) AND 3-
(SUBSTITUTED BENZOTHAZOL-7-YL)-1-SUBSTITUTED-6-
TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONES**

The invention described in this application pertains to weed control in agriculture, horticulture, and other fields where there is a desire to control unwanted plant growth. More specifically, the present application describes certain herbicidal 3-(substituted benzoxazol-7-yl) and 3-(substituted benzothiazol-7-yl)-1-substituted-6-trifluoromethyl-2,4-(1H,3H)pyrimidinediones, as well as compositions containing them, methods of preparing them, and methods for preventing or destroying undesired plant growth by preemergence or postemergence application of herbicidal compositions to the locus where control is desired. The herbicidal activity of the present compounds has not previously been described.

It has now been found that certain 3-(substituted benzoxazol-7-yl) and 3-(substituted benzothiazol-7-yl)-1-substituted-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinediones are highly active herbicides. The novel compounds of the present invention are defined by the following generic structure:



wherein:

R is halogen, straight or branched chain alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkylarylalkyl, haloalkyl, hydroxy, alkoxy, hydroxyalkyl, haloaryl, haloarylalkyl, alkoxyaryl, sulfhydryl, alkylthio, piperidiny, alkylamino, alkoxyalkyl, phenoxy, amino, alkylsulfonylamino, arylsulfonylamino, carboxy, propionyl, halopropionyl, allyloxy, propargyloxy, acetylamino, alkylthienyl,

alkoxyarylalkyl, alkylsulfinyl, alkylsulfonyl, acetoxylalkyl, alkylcarbonyldioxy-alkyl, alkylaminosulfonylamino, haloalkylsulfonylamino, arylaminosulfonylamino, N-alkylsulfonyl-N-alkylamino, N-alkylsulfonyl-N-alkoxyalkylamino, N-alkylsulfonyl-N-alkynylamino, N-alkylsulfonyl-N-alkenylamino, N-arylsulfonyl-
5 N-alkylamino, acetoxylalkylamino, acetoxyl, sodium carboxylate, amino-carboxylate, alkylcarbamoyl, alkylsulfonylcarbamoyl, alkoxy-carbonyl, acetoxylhaloalkyl, acetoxylalkenyl, acetoxylhaloalkenyl, acetoxylalkoxy, alkenyl-oxy, alkynyl-oxy, acetoxylalkylthio, cyanoalkylthio, alkynylthio, arylalkylthio, acetoxylalkylthioalkyl, arylalkylthioalkyl, alkylsulfonylaminoalkyl, alkoxyphos-
10 phinyloxyamino, N-acetyl-N-alkylsulfonylamino, alkynylalkoxyalkyl, alkylcarboxylatephenoxy, halophenoxy, alkylphenoxy, alkoxypropionylphenoxy, or haloalkoxypropionyl;

R¹ is alkyl or amino;

R² is hydrogen or halogen;

15 X is oxygen or sulfur;

Y is hydrogen, halogen, alkoxy, cyano, or nitro, and;

Z is halogen;

where halogen is bromine, chlorine, fluorine, or iodine, and each alkyl moiety has one to six carbon atoms.

20 Preferred compounds are those in which R is alkyl, haloalkyl, piperidinyl, alkylamino, and alkoxyalkyl; R¹ is methyl or amino; R² is hydrogen or chlorine, X is oxygen or sulfur, Y is hydrogen, 4-chloro, 4-bromo, or 4-nitro; and Z is 6-chloro when Y is hydrogen or 6-fluoro when Y is other than hydrogen, where each alkyl moiety has one to four carbon atoms.

25 The compounds of the present invention were prepared by methods known to one skilled in the art.

In general the 3-(substituted-benzoxazol-7-yl)-1-substituted-6-trifluoro-methyl-2,4-(1H,3H)-pyrimidinedione compounds were prepared by one of two routes, depending on whether the benzoxazole ring was formed prior to
30 or after the formation of the 2,4-(1H,3H)-pyrimidinedione ring.

As depicted in Schema 1, those compounds in which the benzoxazole ring was formed prior to the formation of the 2,4-(1H,3H)-pyrimidinedione ring were prepared from one of two starting materials. The first starting material used to prepared those compounds in which the benzoxazole ring is
35 formed first is an hydroxy-, alkyl hydroxy- or alkyl methylhydroxypropionate

(STM1), which is reacted with an alkyl halide, for example methyl iodide, under basic conditions to form the appropriate methoxypropionate (A-1). The propionate (A-1) is in turn hydrogenated under basic conditions to form the corresponding propanoic acid (A-2).

- 5 The second starting material used to prepared those compounds in which the benzoxazole ring is formed first is a substituted-3-chloroaniline (STM2). The aniline (STM2) is reacted with the appropriate alkyl anhydride, for example trimethylacetic anhydride, under acidic conditions, yielding the corresponding N-(substituted-3-chlorophenyl)alkylamide (B-1). The aniline
- 10 (STM2) can also be reacted with the appropriate propanoic acid (A-2), for example 2-methoxy-2-methylpropanoic acid, under basic conditions to form the appropriate N-(substituted-3-chlorophenyl)-substituted-alkylpropionamide (B-2), for example N-(3-chloro-4-fluorophenyl)-2-methoxy-2-methylpropionamide. The amides (B-1 or B-2) are in turn treated with n-butyl-
- 15 lithium and cyclized with solid carbon dioxide, affording the corresponding [2-alkyl- or 2-(substituted-alkyl)-substituted-benzoxazol-7-yl]carboxylic acid (C), for example, (2-t-butyl-6-fluorobenzoxazol-7-yl)carboxylic acid. The so-prepared carboxylic acid (C) is then reacted with ethyl chloroformate, 4-methylmorpholine, and sodium azide, yielding the corresponding [2-alkyl- or
- 20 2-(substituted-alkyl)-substituted-benzoxazol-7-yl]carbonyl azide (C-1), which is treated with ethanol to form the corresponding ethyl N-[2-alkyl- or 2-(substituted-alkyl)-substituted-benzoxazol-7-yl]carbamate (D). Additional substituents may optionally be added to the benzoxazole ring at this point. For example, ethyl N-(2-t-butyl-6-fluorobenzoxazol-7-yl)carbamate (D) is
- 25 treated with N,N-dichlorourethane under acidic conditions, affording ethyl N-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)carbamate (E). The carbamates D and E are then reacted with ethyl 3-amino-4,4,4-trifluorocrotonate, sodium methoxide, and DBU to form the corresponding 3-[2-alkyl- or 2-(substituted-alkyl)-substituted-benzoxazol-7-yl]-6-trifluoromethyl-2,4-(1H,3H)-pyrimidine-
- 30 dione (F). The pyrimidinedione (F) is further reacted under basic conditions with an alkyl halide, for example methyl iodide, or O-(2,4-dinitrophenyl)hydroxylamine, or 1-aminooxysulfonyl-2,4,6-trimethylbenzene, affording the targeted 3-[2-alkyl- or 2-(substituted-alkyl)-substituted-benzoxazol-7-yl]-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (I) or 3-[2-alkyl- or 2-
- 35 (substituted-alkyl)-substituted-benzoxazol-7-yl]-1-amino-6-trifluoromethyl-

2,4-(1H,3H)-pyrimidinedione (II). Examples 1, 2, 4, and 5 provide the detailed procedures for this route.

At this point additional substituents may be optionally added to the pyrimidinedione ring. For example, a chlorine moiety was added to the 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione molecule (I) by treating it with N,N-dichlorourethane under acidic conditions, affording 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-1-methyl-5-chloro-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (III). In addition, the 3-[2-(substituted-methoxyalkyl)-substituted-benzoxazol-7-yl]-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (I), for example 3-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione, can be hydrogenated under basic conditions, yielding the targeted 3-[2-(substituted-hydroxyalkyl)-substituted-benzoxazol-7-yl]-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (IV), for example 3-[2-(1-hydroxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione. Examples 3 and 15 provide the detailed procedures for these methods.

As depicted in Schema 2, those compounds in which the benzoxazole ring was formed after the formation of the 2,4-(1H,3H)-pyrimidinedione ring were prepared by reacting a disubstituted aniline (STM3), for example 4-chloro-2-fluoroaniline, with N,N-dichlorourethane under acid conditions, yielding the appropriate trisubstituted aniline (AA). The anilines (STM3 or AA) are then reacted with pyridine and ethyl chloroformate, yielding the corresponding ethyl N-(disubstituted-or trisubstituted-phenyl)carbamate (H). When the carbamate (H) is an ethyl N-(disubstituted-2-fluorophenyl)carbamate, for example ethyl N-(4-chloro-2,6-difluorophenyl)carbamate, it is nitrated with nitric acid and sulfuric acid to form the corresponding ethyl N-(disubstituted-2-fluoro-3-nitrophenyl)carbamate (H-1). The ethyl N-(disubstituted-2-fluoro-3-nitrophenyl)carbamate (H-1) is in turn reacted with sodium trimethylsilanoate in the presence of methanol and dioxane to form the ethyl N-(disubstituted-2-methoxy-3-nitrophenyl)carbamate (H-2). The carbamate (H-2) is then reduced with iron powder and hydrochloric acid, yielding the corresponding ethyl N-(disubstituted-3-amino-2-methoxyphenyl)carbamate (H-3).

The ethyl N-(disubstituted-phenyl)carbamate (H) is reacted with ethyl 3-amino-4,4,4-trifluorocrotonate, sodium methoxide, and DBU to form the corresponding 3-(disubstituted-phenyl)-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (I), for example, 3-(4-chloro-2-fluorophenyl)-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione. The pyrimidinedione (I) is reacted under basic conditions with an alkyl halide, for example methyl iodide, to form the corresponding 3-(disubstituted-phenyl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (J). The pyrimidinedione (J) is then nitrated with hydrochloric and nitric acid to form the corresponding 3-(2,4-disubstituted-5-nitrophenyl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (K), which is then reduced with iron powder and acetic acid to form the corresponding 3-(5-amino-disubstituted-phenyl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (L). The appropriate carbamate (H-3) is reacted with ethyl 3-amino-4,4,4-trifluorocrotonate, sodium methoxide, and DBU in the manner previously described to form the corresponding 3-(5-amino-6-methoxy-disubstituted-phenyl)-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (LL). The pyrimidinedione (LL) is in turn reacted with an alkyl halide in the manner described previously to form the corresponding 3-(5-amino-6-methoxy-disubstituted-phenyl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (L-1). The pyrimidinedione (L) is reacted with sodium nitrite and sodium azide under acidic conditions, yielding the corresponding 3-(5-azido-2,4-disubstituted-phenyl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (M). The pyrimidinedione (L-1) is treated with boron tribromide and methylene chloride to form the appropriate 3-(5-amino-6-hydroxy-disubstituted-phenyl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (M-1). The pyrimidinediones (M or M-1) are in turn treated with phosphorous pentoxide and hexamethyldisiloxane and then reacted with either an organic acid, for example acetic, chloroacetic, benzoic, substituted benzoic, or methoxyacetic acid, or with the potassium salt of ethyl malonate, affording the targeted 3-(trisubstituted-benzoxazol-7-yl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (V). The appropriate pyrimidinedione (M-1) can also be treated with cyanogen bromide or reacted with potassium ethyl xanthate in the presence of an alcohol to form the targeted 3-(disubstituted-2-amino- or -2-thiobenzoxazol-7-yl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (V), respectively. The pyrimidinedione (M-1) can also be

reacted with 1,1'-carbonyldiimidazole to form the appropriate 3-(disubstituted-2-oxo-2,3-dihydrobenzoxazol-7-yl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (**M-2**). The so-prepared pyrimidinedione (**M-2**) is in turn reacted with phosphorus oxychloride under basic conditions,
5 affording the targeted 3-(disubstituted-2-chlorobenzoxazol-7-yl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (**VI**). The targeted 3-(disubstituted-2-chlorobenzoxazol-7-yl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (**VI**) is in turn reacted with an alcohol under basic conditions, affording the targeted 3-(disubstituted-2-alkoxybenzoxazol-7-yl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (**VII**). Examples 6, 16, 18, 19,
10 24, and 25 provide the detailed procedures for this route.

Additional moieties may also be added to the benzoxazole ring at this point. For example, bromine moieties are added to 3-(2-methyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidine-
15 dione by reacting it with N-bromosuccinimide under basic conditions, yielding the corresponding 3-(2-dibromomethyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (**VB**). An alkyl group can also be added to 3-(2-thio-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione by reacting it with an
20 alkyl halide in the manner described previously, yielding the appropriate 3-(2-alkylthio-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (**VC**). In addition, (**V**), where R in the 2-position is amino may be further reacted to add additional moieties to the amine. For example, an acetyl moiety is added to 3-(2-amino-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione by reacting
25 it with acetyl chloride in the presence of pyridine affording 3-(2-acetylamino-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (**VD**). The so-prepared pyrimidinedione (**VD**) is in turn reacted with sodium hydride and methanesulfonyl chloride to form the
30 appropriate 3-[2-(N-acetyl-N-methylsulfonylamino)-disubstituted-benzoxazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (**VE**), which is hydrogenated under acidic conditions affording 3-(2-methylsulfonylamino-disubstituted-benzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (**VF**). The pyrimidinedione (**VF**) is in turn alkylated in the
35 manner described previously, affording 3-[2-(N-alkyl-N-methylsulfonyl-

amino)-disubstituted-benzoxazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (VG). Examples 14, 17, 20, 21, 22, and 23 provide the detailed procedures for this route.

As depicted in Schema 3, the 3-(substituted-benzothiazol-7-yl)-1-substituted-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinediones are prepared from either of two starting materials, depending upon whether the substituent R in the 2-position of the benzothiazole ring is halogen or alkyl. Those compounds in which the substituent in the 2-position is an alkyl group are prepared by cyclizing a 2,4-, 2,5- or a 2,6-disubstituted-phenylisothiocyanate (STM4) with an alkylmagnesium chloride under basic conditions, yielding the corresponding 2-alkyl-(4- or 6-substituted)benzothiazole (N), for example, 2-methyl-6-fluorobenzothiazole.

Those compounds in which the substituent R in the 2-position of the benzothiazole ring is halogen are prepared by reacting 2-amino-(4- or 6-substituted)benzothiazole (STM5) with t-butyl nitrite and copper(II) chloride to form the corresponding 2-halo-(4- or 6-substituted)benzothiazole (N-1), for example, 2-chloro-6-fluorobenzothiazole. The benzothiazoles (N or N-1) are in turn nitrated with nitric acid and sulfuric acid to form the corresponding 2,4- or 2,6-disubstituted-7-nitrobenzothiazole (O). The so-prepared 7-nitrobenzothiazoles (O) where R in the 2-position of the benzothiazole ring is halogen may then be reacted to replace the 2-halogen with other moieties. For example, 2,4-dichloro-7-nitrobenzothiazole is reacted with piperidine, affording the corresponding 2-(1-piperidinyl)-4- or 6-substituted-7-nitrobenzothiazole (O-1). The 7-nitrobenzothiazoles (O or O-1) are then reduced with iron powder and hydrochloric acid, yielding the corresponding 7-amino-(2,4- or 2,6-disubstituted)benzothiazole (P).

At this point the 7-aminobenzothiazole (P), where R in the 2-position is halogen, alkyl, or other, such as 1-piperidinyl, may be treated directly with ethyl chloroformate to form the corresponding ethyl benzothiazolylcarbamate (Q), or with phosgene to form the corresponding benzothiazolylisocyanate (R-1). In an alternate method, (P), where R in the 2-position is halogen, may be reacted to replace the halogen with other moieties, or to add additional moieties to the benzothiazole ring. For example, (P), where R in the 2-position is chlorine, is reacted with sodium thiomethoxide, yielding the corresponding 7-amino-2-methylthio-(4- or 6-substituted)benzothiazole (P-

1); or (P), where R in the 2-position is chlorine, is reacted with N-chlorosuccinimide, yielding the corresponding 7-amino-2,4,6-trichlorobenzothiazole (P-2). The 7-aminobenzothiazoles (P-1) prepared as outlined above may now be reacted with ethyl chloroformate to form the corresponding ethyl benzothiazolylcarbamates (Q), or the 7-aminobenzothiazoles (P-2) may be reacted with phosgene to form the corresponding benzothiazolylisocyanates (R-1). Examples 9 and 10 provided the detailed procedures for these methods.


Additional moieties may also be added to the ethyl benzothiazolylcarbamate (Q) at this point. For example, a chlorine moiety is added to ethyl (2-chloro- or 2-methyl-6-fluorobenzothiazol-7-yl)carbamate by treating it with N,N-dichlorourethane under acidic conditions, yielding the corresponding ethyl (2-substituted-4-chloro-6-fluorobenzothiazol-7-yl)carbamate (Q-1). In addition, (Q-1), where R in the 2-position is chlorine, may be further reacted to substitute other moieties in the 2-position. For example, ethyl (2, 4-dichloro-6-fluorobenzothiazol-7-yl)carbamate (Q-1) is reacted with 2-methylpropylamine, affording ethyl (2-methylpropyl-4-chloro-6-fluorobenzothiazol-7-yl)carbamate (Q-2). Examples 11, 12, and 13 provide the detailed procedures for these methods.

By the methods previously described, the ethyl benzothiazolylcarbamates (Q, Q-1, or Q-2) and the benzothiazolylisocyanates (R-1) are then cyclized to the corresponding pyrimidinediones (S) with ethyl 3-amino-4,4,4-trifluorocrotonate in the presence of DBU, which are in turn alkylated, yielding the targeted 3-(2,4- or 2,6- or 2,4,6-substituted-benzothiazol-7-yl)-1-alkyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (VIII). In one case alkylation of (S) resulted not only in alkylation at the expected 1-position of the pyrimidinedione ring, but also replaced the chlorine in the 4-position of the benzothiazole ring with methoxy. Examples 7 and 10 provide the detailed procedures for this route.


Once the pyrimidinedione ring (VIII) is formed, additional substituents may be optionally added. For example, a chlorine moiety is added to the 3-(2-chloro-6-fluorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione molecule by treating it with N-chlorosuccinimide under acidic conditions, affording 3-(2-chloro-6-fluorobenzothiazol-7-yl)-1-methyl-5-

chloro-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (IX). Example 8 provides the detailed procedure for this method.



Schema 1




Schema 1 (continued)




Schema 1 (cont)




Schema 2



Schema 2 (continued)







Schema 2 (continued)




16


Schema 2 (continued)








Schema 2 (continued)




Schema 3



Schema 3 (cont)



Schema 3 (cont)



EXAMPLE 1

SYNTHESIS OF 3-(2-t-BUTYL-4-CHLORO-6-FLUOROBENZOXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE
(COMPOUND 4)5 Step A Synthesis of N-(3-chloro-4-fluorophenyl)-t-butylamide as an
 intermediate

 A solution of 10 grams (69 mmole) of 3-chloro-4-fluoroaniline, 75 mL of acetic acid, and 10 mL of tetrahydrofuran was stirred and cooled in an ice bath. During 15 minutes, 20 mL of trimethylacetic anhydride was added
10 dropwise. Upon completion of the addition the reaction mixture was allowed to warm to ambient temperature, where it stirred for six hours. The reaction mixture was then poured into 300 mL of ice-water and stirred for 15 minutes. The resulting precipitate was collected by filtration and washed with water. The precipitate was purified by column chromatography on silica gel with 4:1
15 hexane:ethyl acetate as eluant. The product-containing fractions were combined and concentrated under reduced pressure, yielding 10 grams of N-(3-chloro-4-fluorophenyl)-t-butylamide. The NMR spectrum was consistent with the proposed structure.

20 Step B Synthesis of (2-t-butyl-6-fluorobenzoxazol-7-yl)carboxylic acid
 as an intermediate

 Under a nitrogen atmosphere a stirred solution of 5.0 grams (22 mmole) of N-(3-chloro-4-fluorophenyl)-t-butylamide in 100 mL of tetrahydrofuran was cooled to about -50 °C, and 18 mL (45 mmole) of 2.5 M n-butyllithium in hexanes was added dropwise at a rate to maintain the reaction mixture
25 temperature below -30 °C. Upon completion of the addition the reaction mixture was stirred for 90 minutes and poured over solid carbon dioxide. The mixture was allowed to warm to ambient temperature, and 150 mL of aqueous 1 N sodium hydroxide solution was added. The mixture was extracted with two 50 mL portions of ethyl acetate. The organic extracts
30 were combined and washed with 50 mL of aqueous 1 N sodium hydroxide solution. The combined washes were acidified to a pH of 2 with concentrated hydrochloric acid and extracted with three 50 mL portions of ethyl acetate. The extracts were combined and washed with aqueous saturated sodium chloride solution. The organic layer was dried with magnesium
35 sulfate and filtered. The filtrate was concentrated under reduced pressure,

yielding 3.8 grams of (2-t-butyl-6-fluorobenzoxazol-7-yl)carboxylic acid. The NMR spectrum was consistent with the proposed structure.

Step C Synthesis of ethyl N-(2-t-butyl-6-fluorobenzoxazol-7-yl) carbamate as an intermediate

5 Under a nitrogen atmosphere a stirred solution of 3.7 grams (15.5 mmole) of (2-t-butyl-6-fluorobenzoxazol-7-yl)carboxylic acid in 40 mL of tetrahydrofuran was cooled in an ice bath, and 1.6 grams (16 mmole) of 4-methylmorpholine was added slowly. To this was then added 1.5 mL (16 mmole) of ethyl chloroformate during a five minute period. Upon completion
10 of the addition the reaction mixture was stirred for 30 minutes, after which a solution of 2.0 grams (32 mmole) of sodium azide in 20 mL of water was added dropwise during a 20 minute period. The reaction mixture was allowed to warm to ambient temperature as it stirred for 30 minutes. The reaction mixture was then analyzed by thin layer chromatography (TLC),
15 which indicated that no starting acid remained. The reaction mixture was poured into 150 mL of aqueous 5% hydrochloric acid solution, and the mixture was extracted with diethyl ether. The combined extract was washed with an aqueous saturated sodium chloride solution, dried with magnesium sulfate, and filtered. The filtrate was concentrated under reduced pressure,
20 yielding 3.4 grams of the azide intermediate as a yellow oil, which was taken up in 50 mL of ethanol and heated at reflux for 18 hours. After this time TLC analysis indicated that the reaction to the carbamate was complete. The reaction mixture was concentrated under reduced pressure, taken up in 50 mL of aqueous 1 N hydrochloric acid solution, and extracted with three 50 mL portions of diethyl ether. The combined extracts were washed with an
25 aqueous saturated sodium chloride solution, dried with magnesium sulfate, and filtered. The filtrate was concentrated under reduced pressure, yielding 3.2 grams of ethyl N-(2-t-butyl-6-fluorobenzoxazol-7-yl)carbamate. The NMR spectrum was consistent with the proposed structure.

30 **Step D Synthesis of ethyl N-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)carbamate as an intermediate**

Under a nitrogen atmosphere, a solution of 3.1 grams (11.0 mmole) of ethyl N-(2-t-butyl-6-fluorobenzoxazol-7-yl)carbamate, 35 mL of acetic acid, and 0.2 mL (2.4 mmole) of concentrated hydrochloric acid was stirred and
35 cooled in an ice bath. During five minutes 2.2 grams (14 mmole) of N,N-

dichlorourethane was added via syringe. Upon completion of the addition the reaction mixture was stirred for 4 hours, then analyzed by TLC, which indicated the reaction was complete. The reaction mixture was poured into 100 mL of water, and 15 mL of aqueous saturated sodium bicarbonate solution was carefully added. The reaction mixture was then extracted with two 75 mL portions of diethyl ether, and the combined extracts were concentrated under reduced pressure, yielding a residual oil. The oil was subjected to column chromatography on silica gel with 8:1 hexane:ethyl acetate as the eluant. The product-containing fractions were combined and concentrated under reduced pressure, yielding 2.4 grams of ethyl N-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)carbamate, m.p. 127-129 °C. The NMR spectrum was consistent with the proposed structure.

Step E Synthesis of 3-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

Under a nitrogen atmosphere a solution of 0.33 gram (6.2 mmole) of sodium methoxide in 45 mL of N,N-dimethylformamide was stirred and cooled in an ice bath. During 10 minutes a solution of 1.14 grams (6.2 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate in 10 mL of N,N-dimethylformamide was added dropwise. Upon completion of the addition the reaction mixture was stirred for 30 minutes, after which a solution of 1.8 grams (5.9 mmole) of ethyl N-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)carbamate in 30 mL of N,N-dimethylformamide was added dropwise during a 15 minute period. The reaction mixture was heated to 50 °C, and 0.91 gram (6 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene was added via pipette. Upon completion of the addition the reaction mixture was heated at 120-125 °C for 24 hours, then cooled to ambient temperature, where it stirred for 24 hours. After this time the reaction mixture was poured into 100 mL of aqueous 10% lithium chloride solution and acidified with concentrated hydrochloric acid. The mixture was then extracted with three 50 mL portions of ethyl acetate. The combined extracts were concentrated under reduced pressure, yielding a black oil, which was extracted with four 50 mL portions of aqueous saturated sodium bicarbonate solution. The combined extracts were acidified with concentrated hydrochloric acid and then extracted with three 40 mL portions of ethyl acetate. The combined extracts were dried with magnesium sulfate and filtered. The filtrate was concentrated under

reduced pressure, yielding 0.34 gram of a dark oil. The black oil collected earlier was purified by column chromatography on silica gel. Elution was accomplished with 3:1 hexane:ethyl acetate. The product-containing fractions were combined and concentrated under reduced pressure, yielding 5 0.65 gram of a brown solid. The brown solid and the dark oil were combined, yielding 3-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione. The NMR spectrum was consistent with the proposed structure.

An alternative method used for preparing the 3-(2-t-butyl-4-chloro-6-10 fluorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione intermediate is as follows. Under a nitrogen atmosphere a solution of 2.4 grams (45 mmole) of sodium methoxide in 40 mL of N,N-dimethylformamide was stirred and cooled in an ice bath. During 15 minutes a solution of 8.2 grams (45 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate in 40 mL of N,N-15 dimethylformamide was added dropwise. Upon completion of the addition the reaction mixture was stirred at 10 °C for 1.5 hours, after which a solution of 13.0 grams (41 mmole) of ethyl N-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)carbamate (prepared in the manner of Steps A-D, Example 1) in 40 mL of N,N-dimethylformamide was added dropwise during a 15 minute period. 20 Upon completion of the addition the reaction mixture was warmed to ambient temperature, where it stirred for five minutes, then was heated to 60 °C, and 6.8 grams (45 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene was added via pipette. Upon completion of the addition, the reaction mixture was heated at 120 °C for 4 hours, then cooled to ambient temperature, where it stirred for 25 about 18 hours. The reaction mixture was poured into 200 mL of aqueous 10% potassium carbonate, 100 mL of water was added, and the mixture was extracted with two 100 mL portions of diethyl ether. The combined extracts were washed with 50 mL of aqueous saturated sodium bicarbonate solution. The combined aqueous layer and washes were acidified with about 30 mL of 30 concentrated hydrochloric acid, and the mixture was extracted with three 100 mL portions of ethyl acetate. The combined organic extracts were washed with two 50 mL portions of aqueous saturated sodium chloride solution and two 50 mL portions of aqueous 10% lithium chloride solution. The organic layer was dried with magnesium sulfate, filtered, and concentrated under 35 reduced pressure, yielding 6.4 grams of 3-(2-t-butyl-4-chloro-6-

fluorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, a dark oil. The NMR spectrum was consistent with the proposed structure.

Step F Synthesis of 3-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione
(Compound 4)

5 A stirred solution of 0.85 gram (2.1 mmole) of 3-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 0.48 gram (3.5 mmole) of potassium carbonate, 0.91 gram (6.4 mmole) of iodo-
10 methane, and 35 mL of acetone was heated at 40 °C for 2 hours, then stirred at ambient temperature for 20 hours. The reaction mixture was concentrated under reduced pressure, and the concentrate was extracted with two 50 mL portions of diethyl ether. The combined extracts were dried with magnesium sulfate and filtered. The filtrate was concentrated under
15 reduced pressure to a yellow oil, which was subjected to column chromatography on silica gel. Elution was accomplished with 8:1 hexane:ethyl acetate. The product-containing fractions were combined and concentrated under reduce pressure, yielding 0.60 gram of 3-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, m.p. 68-70 °C. The NMR spectrum was consistent with the proposed structure.

20 An alternative method for preparing 3-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione is the following: A solution of 16.0 grams (39 mmole) of 3-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 8.3 grams (60 mmole) of potassium carbonate, 8.5 grams (60 mmole) of iodo-
25 methane, and 150 mL of acetone was stirred at ambient temperature for about 18 hours. The reaction mixture was then concentrated under reduced pressure, and the residue was taken up in aqueous 5% hydrochloric acid. The mixture was extracted with three 75 mL portions of diethyl ether. The combined extracts were washed with aqueous saturated sodium chloride
30 solution. The organic layer was dried with magnesium sulfate, filtered, and concentrated under reduced pressure, yielding 12.7 grams of a dark oil, which was subjected to column chromatography on silica gel. Elution was accomplished with 6:1 heptane:ethyl acetate. The product-containing fractions were combined and concentrated under reduced pressure, yielding
35 9.4 grams of 3-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-

trifluoromethyl-2,4-(1H,3H)pyrimidinedione. The NMR spectrum was consistent with the proposed structure.

EXAMPLE 2

5 SYNTHESIS OF 3-(2-t-BUTYL-4-CHLOROBENZOXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE (COMPOUND 1)

Step A Synthesis of N-(2,5-dichlorophenyl)-t-butylamide as an intermediate

10 Under a nitrogen atmosphere a stirred solution of 5.0 grams (31 mmole) of 2,5-dichloroaniline, 6.5 grams (35 mmole) of trimethylacetic anhydride, and 1 drop of concentrated sulfuric acid was heated at about 80°C for four hours, after which analysis by thin layer chromatography (TLC) indicated that the reaction was complete. The reaction mixture was poured into 150 mL of
15 aqueous 1 N hydrochloric acid and stirred for 30 minutes. The resulting precipitate was collected by filtration and washed with water. The filtrate and washes were combined and dried in a vacuum oven at 45 °C for about 16 hours, yielding 7.3 grams of N-(2,5-dichlorophenyl)-t-butylamide. The NMR spectrum was consistent with the proposed structure.

20 Step B Synthesis of (2-t-butyl-4-chlorobenzoxazol-7-yl)carboxylic acid as an intermediate

 This compound was prepared in the manner of Step B, Example 1, with 6.0 grams (24 mmole) of N-(2,5-dichlorophenyl)-t-butylamide, 100 mL of tetrahydrofuran, and 24 mL (60 mmole) of 2.5 M n-butyllithium in hexanes
25 as reagents. The yield of (2-t-butyl-4-chlorobenzoxazol-7-yl)carboxylic acid was 4.2 grams. The NMR spectrum was consistent with the proposed structure.

Step C Synthesis of ethyl N-(2-t-butyl-4-chlorobenzoxazol-7-yl)carbamate as an intermediate

30 This compound was prepared in the manner of Step C, Example 1, with 3.0 grams (11.8 mmole) of (2-t-butyl-4-chlorobenzoxazol-7-yl)carboxylic acid, 40 mL of tetrahydrofuran, 1.21 grams (12 mmole) of 4-methylmorpholine, and 1.3 grams (12 mmole) of ethyl chloroformate as reagents. The yield of ethyl N-(2-t-butyl-4-chlorobenzoxazol-7-yl)carbamate was 2.9 grams. The
35 NMR spectrum was consistent with the proposed structure.

Step D Synthesis of 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

This compound was prepared in the manner of Step E, Example 1, with 0.65 gram (12.0 mmole) of sodium methoxide, 45 mL of N,N-dimethyl-
5 formamide, 2.0 grams (11.0 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate, 2.9 grams (9.8 mmole) of ethyl N-(2-t-butyl-4-chlorobenzoxazol-7-yl)carbamate, and 1.7 grams (11 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene as reagents. The yield of 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 2.8 grams. The NMR
10 spectrum was consistent with the proposed structure.

Step E Synthesis of 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione (Compound 1)

This compound was prepared in the manner of Step F, Example 1, with 2.0 grams (5.2 mmole) of 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 1.1 grams (7.8 mmole) of potassium
15 carbonate, 1.1 grams (8 mmole) of iodomethane, and 50 mL of acetone as reagents. The yield of 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, m.p. 146-148 °C, was 1.5 grams. The NMR spectrum was consistent with the proposed structure.

20

EXAMPLE 3

SYNTHESIS OF 3-(2-t-BUTYL-4-CHLOROBENZOXAZOL-7-YL)-1-METHYL-5-CHLORO-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE (COMPOUND 2)

25 A stirred solution of 0.84 gram (2.1 mmole) of 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione (prepared as described in Example 2) in 20 mL of acetic acid had 0.16 gram (1.0 mmole) of N,N-dichlorourethane added via syringe. Upon completion of the addition the clear reaction mixture was stirred for 15 minutes, then one drop
30 of concentrated hydrochloric acid was added, and the reaction mixture was stirred at ambient temperature for 72 hours. After this time the reaction mixture was poured into 100 mL of water, and 10 grams of sodium carbonate were added. The mixture was extracted with three 50 mL portions of methylene chloride. The organic extracts were combined and washed
35 with one 50 mL portion of an aqueous 5% sodium carbonate solution. The

organic layer was dried with sodium sulfate and filtered. The filtrate was concentrate under reduced pressure, yielding a pale yellow oil which was subjected to column chromatography on silica gel. Elution was accomplished with 9:1 heptane:ethyl acetate, followed by 2:1 heptane:ethyl acetate. The product-containing fractions were combined and concentrated under reduced pressure, yielding 0.57 gram of 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-1-methyl-5-chloro-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, m.p. 83-86 °C. The NMR spectrum was consistent with the proposed structure.

10

EXAMPLE 4

SYNTHESIS OF 3-(2-t-BUTYL-4-CHLOROBENZOXAZOL-7-YL)-1-AMINO-6-TRIFLUOROMETHYL-2,4-(1H,3H)-PYRIMIDINEDIONE (COMPOUND 3)

15 A stirred solution of 0.5 gram (1.3 mmole) of 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione (prepared in Step D, Example 2), 0.21 gram (1.5 mmole) of potassium carbonate, and 30 mL of N,N-dimethylformamide was heated at 60 °C for 20 minutes, cooled to ambient temperature, and 0.30 gram (1.5 mmole) of O-(2,4-dinitrophenyl)-
20 hydroxylamine was added. Upon completion of the addition the reaction mixture was stirred for about 18 hours, then poured into 100 mL of water and extracted with three 40 mL portions of methylene chloride. The combined organic extracts were dried with magnesium sulfate, filtered, and concentrated under reduced pressure, yielding 0.3 gram of a yellow oil. The
25 aqueous layer was neutralized with excess ammonium chloride and extracted with two 50 mL portions of ethyl acetate. The combined extracts were washed with an aqueous saturated sodium chloride solution. The organic layer was dried with magnesium sulfate, filtered, and concentrated under reduced pressure, yielding 0.22 gram of a solid. Thin layer chromatography (TLC) indicated that this solid was the starting material, 3-(2-t-butyl-
30 4-chlorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione. The yellow oil was subjected to column chromatography on silica gel. Elution was accomplished with 5:1 heptane:ethyl acetate. The product-containing fractions were combined and concentrated under reduced pressure, yielding
35 0.12 gram of 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-1-amino-6-trifluoromethyl-

2,4-(1H,3H)pyrimidinedione, m.p. 104-106°C. A solution containing the recovered 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 1.1 grams of potassium carbonate, 10 mL of N,N-dimethylformamide, and 1.1 grams of O-(2,4-dinitrophenyl)hydroxylamine
5 was stirred for about 60 hours. After this time the work-up procedure described above was repeated, yielding 0.7 gram of solid. Analysis by TLC indicated that the solid was 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-1-amino-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, m.p. 104-106 °C. The two
10 materials were combined for a yield of 0.8 gram of 3-(2-t-butyl-4-chlorobenzoxazol-7-yl)-1-amino-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione. The NMR spectrum was consistent with the proposed structure.

EXAMPLE 5

15 SYNTHESIS OF 3-(2-t-BUTYL-4-CHLORO-6-FLUOROBENZOXAZOL-7-YL)-1-AMINO-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE (COMPOUND 5)

Step A Synthesis of 1-aminoxysulfonyl-2,4,6-trimethylbenzene as an intermediate

To a stirred solution of 25.0 grams (114 mmole) of 2,4,6-trimethyl-
20 benzenesulfonyl chloride, 15.2 grams (114 mmole) of t-butyl-N-hydroxycarbamate, and 350 mL of diethyl ether, cooled in an ice bath, was added dropwise 11.4 grams (114 mmole) of triethylamine. Upon completion of the addition the reaction mixture was stirred at ambient temperature for about 18 hours, then filtered, and the filtrate was concentrated under reduced
25 pressure. The concentrate was taken up in toluene and petroleum ether, filtered, and 150 mL of trifluoroacetic acid was added. The reaction mixture was stirred for 1.5 hours in an ice bath, then poured into ice-water, filtered, and extracted with diethyl ether. The organic extract was dried with
30 magnesium sulfate and filtered. Petroleum ether was added to the filtrate, and the resulting precipitate was collected by filtration, yielding 14.4 grams of 1-aminoxysulfonyl-2,4,6-trimethylbenzene. The NMR spectrum was consistent with the proposed structure.

Step B Synthesis of 3-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-amino-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione
35 (Compound 5)

Under a nitrogen atmosphere a stirred solution of 1.0 gram (2.5 mmole) of 3-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione (prepared in Example 1, Steps A-E), 0.41 gram (3.0 mmole) of potassium carbonate, and 50 mL of tetrahydrofuran was cooled in an ice bath. To this was added 1.1 grams (about 5 mmole) of 1-amino-oxysulfonyl-2,4,6-trimethylbenzene. Upon completion of the addition the reaction mixture was stirred in the ice bath for ten minutes, then warmed to ambient temperature, where it stirred for an additional 18 hours. After this time the reaction mixture was poured into 150 mL of aqueous 10% ammonium chloride solution and extracted with two 75 mL portions of diethyl ether. The combined extracts were washed with two 50 mL portions of an aqueous saturated sodium chloride solution. The organic layer was dried with magnesium sulfate and filtered. The filtrate was concentrated under reduced pressure to an oil, which was subjected to column chromatography on silica gel. Elution was accomplished with 6:1 heptane:ethyl acetate. The product-containing fractions were combined and concentrated under reduced pressure, yielding 0.75 gram of 3-(2-t-butyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-amino-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, m.p. 90-92 °C. The NMR spectrum was consistent with the proposed structure.

EXAMPLE 6

SYNTHESIS OF 3-(2-METHYL-4-CHLORO-6-FLUOROBENZOXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE (COMPOUND 6)

Step A Synthesis of ethyl N-(4-chloro-2-fluorophenyl)carbamate as an intermediate

A solution of 6.9 grams (47 mmole) of 4-chloro-2-fluoroaniline in 75 mL of pyridine was stirred and cooled in an ice bath. During 15 minutes, 5.4 grams (50 mmole) of ethyl chloroformate was added dropwise. Upon completion of the addition the reaction mixture was heated at 60-70 °C for 1.5 hours, then concentrated under reduced pressure to a residue. The residue was stirred with 150 mL of aqueous 3 N hydrochloric acid, and the resulting precipitate was collected by filtration. The precipitate was washed with aqueous 3 N hydrochloric acid and then with water. The precipitate was dried under ambient conditions for about 18 hours, then dried under vacuum

for two hours, yielding 9.6 grams of ethyl N-(4-chloro-2-fluorophenyl)carbamate. The NMR spectrum was consistent with the proposed structure.

Step B Synthesis of 3-(4-chloro-2-fluorophenyl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

5 This compound was prepared in a manner analogous to Step E, Example 1 with 1.1 grams (20 mmole) of sodium methoxide, 45 mL of N,N-dimethylformamide, 3.7 grams (20 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate, 3.8 grams (17.5 mmole) of ethyl N-(4-chloro-2-fluorophenyl)carbamate, and 3.0 grams (20 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene as reagents. This reaction differed in that the reaction mixture was heated at 105 °C for four hours rather than 120-125 °C for 24 hours. The yield of 3-(4-chloro-2-fluorophenyl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 4.0 grams. The NMR spectrum was consistent with the proposed structure.

15 Step C Synthesis of 3-(4-chloro-2-fluorophenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

 This compound was prepared in the manner of Step F, Example 1 with 4.0 grams (13.2 mmole) of 3-(4-chloro-2-fluorophenyl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 4.2 grams (30 mmole) of potassium carbonate, 4.3 grams (30 mmole) of iodomethane, and 55 mL of acetone as reagents. The yield of 3-(4-chloro-2-fluorophenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 3.0 grams. The NMR spectrum was consistent with the proposed structure.

25 Step D Synthesis of 3-(4-chloro-2-fluoro-5-nitrophenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

 A solution of 3.0 grams (9.4 mmole) of 3-(4-chloro-2-fluorophenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione in 30 mL of concentrated sulfuric acid was stirred and cooled in an ice bath. During 15 minutes, 0.75 mL (12 mmole) of aqueous 70% nitric acid was added dropwise, after which the reaction mixture was allowed to warm to ambient temperature, where it stirred for two hours. The reaction mixture was then poured into 150 mL of ice-water. The resulting precipitate was collected by filtration, washed with water, and dried under reduced pressure, yielding 3.0 grams of 3-(4-chloro-2-fluoro-5-nitrophenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-

pyrimidinedione. The NMR spectrum was consistent with the proposed structure.

Step E Synthesis of 3-(5-amino-4-chloro-2-fluorophenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

5 Under a nitrogen atmosphere a solution of 3.0 grams (8.2 mmole) of 3-(4-chloro-2-fluoro-5-nitrophenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione, and 20 mL of acetic acid in 80 mL of ethanol was stirred, and 1.8 grams (32 mmole) of iron powder was added. The reaction mixture
10 was then heated under reflux for one hour, after which analysis by thin layer chromatography (TLC) indicated that the reaction was complete. The reaction mixture was concentrated under reduced pressure. The concentrate was then taken up in 50 mL of aqueous 5% sodium bicarbonate solution and 100 mL of ethyl acetate. The mixture was filtered through
15 diatomaceous earth, and the filter cake was washed with 20 mL of water and three 50 mL portions of ethyl acetate. The filtrate was placed in a separatory funnel and 100 mL of aqueous 5% sodium bicarbonate solution was added. The aqueous phase was separated and the organic phase was washed with
20 50 mL of aqueous saturated sodium bicarbonate solution, then with 50 mL of aqueous saturated sodium chloride solution. The organic layer was dried with magnesium sulfate and filtered. The filtrate was concentrated under reduced pressure, yielding 2.6 grams of 3-(5-amino-4-chloro-2-fluorophenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione. The NMR spectrum was consistent with the proposed structure.

25 Step F Synthesis of 3-(5-azido-4-chloro-2-fluorophenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

 A stirred solution of 6.0 mL of concentrated sulfuric acid and 15 mL of water was cooled to 5 °C in an ice bath. To this was added 1.6 grams (4.7 mmole) of 3-(5-amino-4-chloro-2-fluorophenyl)-1-methyl-6-trifluoromethyl-
30 2,4-(1H,3H)pyrimidinedione. Over a period of 10 minutes a solution of 0.36 gram (5.2 mmole) of sodium nitrite in 5.0 mL of water was added dropwise at a rate to maintain the reaction mixture temperature below 10 °C. Upon completion of the addition the reaction mixture was stirred for 40 minutes at about 5 to 10 °C. A solution of 0.37 gram (5.6 mmole) of sodium azide in 5.0
35 mL of water was then added during a 20 minute period at a rate to maintain

the reaction mixture temperature below 10 °C. The reaction mixture was then stirred for 20 minutes and poured into 150 mL of water. The resulting precipitate was filtered, washed with water, and dried at ambient conditions for about 18 hours, then at 65 °C for 6 hours under vacuum. The yield of 3-(5-azido-4-chloro-2-fluorophenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was about 1.8 grams. The NMR spectrum was consistent with the proposed structure.

Step G Synthesis of 3-(2-methyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione
(Compound 6)

Under a nitrogen atmosphere a stirred solution of 4.0 grams (28 mmole) of phosphorus pentoxide and 10 mL (47 mmole) of hexamethyldisiloxane was heated at 90 °C for 30 minutes. To this was added dropwise during a 5.0 minute period a solution of 0.80 gram (2.2 mmole) of 3-(5-azido-4-chloro-2-fluorophenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione in 5.0 mL of acetic acid. Upon completion of the addition the reaction mixture was heated at reflux for 5.0 hours, cooled to ambient temperature, and poured into 50 mL of water. The reaction mixture was extracted with three 25 mL portions of ethyl acetate. The combined extracts were washed with 50 mL of aqueous saturated sodium bicarbonate solution and 50 mL of aqueous saturated sodium chloride solution. The organic layer was dried with magnesium sulfate and filtered. The filtrate was concentrated under reduced pressure, yielding 0.7 gram of a residual oil, which was subjected to column chromatography on silica gel. Elution was accomplished with 3:1 heptane:ethyl acetate. The product-containing fractions were combined and concentrated under reduced pressure, yielding 0.40 gram of an oily white solid, 3-(2-methyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, m.p. 64-66 °C. The NMR spectrum was consistent with the proposed structure.

EXAMPLE 7

SYNTHESIS OF 3-(2-CHLORO-6-FLUOROBENZOTHAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE
(COMPOUND 10)

Step A Synthesis of 2-chloro-6-fluoro-7-nitrobenzothiazole as an intermediate

5 This compound was prepared in the manner of Step D, Example 6, with 5.0 grams (26.6 mmole) of 2-chloro-6-fluorobenzothiazole, 20 mL of concentrated sulfuric acid, and 2.0 mL (35 mmole) of concentrated nitric acid as reagents. This procedure differed in that the reaction mixture was neutralized with 20 mL of concentrated ammonium hydroxide before the precipitate was filtered. The yield of 2-chloro-6-fluoro-7-nitrobenzothiazole was 4.4 grams. The NMR spectrum was consistent with the proposed structure.

10 **Step B Synthesis of 7-amino-2-chloro-6-fluorobenzothiazole as an intermediate**

15 This compound was prepared in the manner of Step E, Example 6, with 4.2 grams (18 mmole) of 2-chloro-6-fluoro-7-nitrobenzothiazole, 4.0 grams (72 mmole) of iron powder, 10 mL of acetic acid, 100 mL of ethanol, and 2 mL (24 mmole) of concentrated hydrochloric acid as reagents. The yield of 7-amino-2-chloro-6-fluorobenzothiazole was 2.7 grams. The NMR spectrum was consistent with the proposed structure.

Step C Synthesis of ethyl N-(2-chloro-6-fluorobenzothiazol-7-yl) carbamate as an intermediate

20 Under a nitrogen atmosphere 20 mL of pyridine was stirred, and 2.7 grams (25 mmole) of ethyl chloroformate was added via syringe. Upon completion of the addition the reaction mixture was stirred for 15 minutes. To this was then added a solution of 2.6 grams (12.8 mmole) of 7-amino-2-chloro-6-fluorobenzothiazole in 30 mL pyridine dropwise during a 15 minute period. The reaction mixture was warmed to ambient temperature, where it stirred for one hour, after which the reaction mixture was poured into 200 mL of 1 N aqueous hydrochloric acid. The resulting precipitate was collected by filtration, washed with water, and dried under reduced pressure. The yield of ethyl N-(2-chloro-6-fluorobenzothiazol-7-yl)carbamate was 2.8 grams. The NMR spectrum was consistent with the proposed structure.

30 **Step D Synthesis of 3-(2-chloro-6-fluorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate**

35 This compound was prepared in the manner of Step E, Example 1, with 2.8 grams (10 mmole) of ethyl N-(2-chloro-6-fluorobenzothiazol-7-yl)carbamate, 2.2 grams (12 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate, 0.6

gram (11 mmole) of sodium methoxide, 30 mL of N,N-dimethylformamide, and 1 mL (7.3 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene as reagents. The yield of 3-(2-chloro-6-fluorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 2.2 grams. The NMR spectrum was consistent with the proposed structure.

Step E Synthesis of 3-(2-chloro-6-fluorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione (Compound 10)

This compound was prepared in the manner of Step F, the alternate method, Example 1, with 2.2 grams (6.0 mmole) of 3-(2-chloro-6-fluorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 1.4 grams (10 mmole) of potassium carbonate, 1.4 grams (10 mmole) of iodomethane, and 50 mL of acetone as reagents. The yield of 3-(2-chloro-6-fluorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 1.3 grams, m.p. 69-71 °C. The NMR spectrum was consistent with the proposed structure.

EXAMPLE 8

SYNTHESIS OF 3-(2-CHLORO-6-FLUOROBENZOTHAZOL-7-YL)-1-METHYL-5-CHLORO-6-TRIFLUOROMETHYL-2,4-(1H,3H)-PYRIMIDINEDIONE (COMPOUND 11)

Under a nitrogen atmosphere a stirred solution of 0.6 gram of (1.5 mmole) of 3-(2-chloro-6-fluorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione (prepared by way of Example 7), 0.24 gram (1.8 mmole) of N-chlorosuccinimide, 15 mL of N,N-dimethylformamide, and 0.1 mL acetic acid was heated at 65 °C for 24 hours. After this time the reaction mixture was cooled to ambient temperature and poured into 150 mL of aqueous 5% sodium bicarbonate. To this was added 10 grams of solid lithium chloride. The reaction mixture was then stirred for ten minutes. The resulting precipitate was collected by filtration and dried to give 0.42 gram of a white solid, which was purified by column chromatography on silica gel. Elution was accomplished using 6:1 hexane:ethyl acetate. The product-containing fractions were combined and concentrated under reduced pressure, yielding 0.26 gram of 3-(2-chloro-6-fluorobenzothiazol-7-yl)-1-methyl-5-chloro-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, m.p. 81-83 °C. The NMR spectrum indicated that the material was 3-(2-chloro-6-fluoro-

benzothiazol-7-yl)-1-methyl-5-chloro-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione rather than 3-(2,4-dichloro-6-fluorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione.

EXAMPLE 9

**SYNTHESIS OF 3-(2-METHYLTHIO-4-CHLOROBENZOTHAZOL-7-
YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)-
PYRIMIDINEDIONE (COMPOUND 8)**

Step A **Synthesis of 7-amino-2-methylthio-4-chlorobenzothiazole as an intermediate**

Under a nitrogen atmosphere a stirred solution of 0.88 gram (4 mmole) of 7-amino-2,4-dichlorobenzothiazole, 1.2 grams (17 mmole) of sodium thiomethoxide, and 25 mL of N,N-dimethylformamide was heated at 60 °C for two hours. The reaction mixture was cooled, poured into 150 mL of cold 5% sodium bicarbonate, and 50 mL of saturated sodium chloride solution was added. The resulting precipitate was filtered and vacuum dried, yielding 0.8 grams of a yellow green solid. NMR analysis indicated this material was half 7-amino-2,4-dichlorobenzothiazole starting material and half the desired product 7-amino-2-methylthio-4-chlorobenzothiazole.

The reaction was repeated with 2.3 grams (10.5 mmole) of 7-amino-2,4-dichlorobenzothiazole, 1.2 grams (17 mmole) of sodium thiomethoxide, and 25 mL of N,N-dimethylformamide as reagents. After the reaction mixture had been heated for two hours at 60 °C the 0.8 grams from the previous reaction was added, and the reaction mixture was heated for an additional three hours. The reaction mixture was cooled and poured into 300 mL of cold aqueous sodium bicarbonate solution. To this was then added 10 grams of solid lithium chloride. The reaction mixture was extracted with three 100 mL portions of diethyl ether. The combined organic extracts were washed with an aqueous saturated sodium chloride solution. The organic layer was dried with sodium sulfate, filtered, and concentrated under reduced pressure, yielding 2.0 grams of 7-amino-2-methylthio-4-chlorobenzothiazole. The NMR spectrum was consistent with the proposed structure.

Step B **Synthesis of ethyl N-(2-methylthio-4-chlorobenzothiazol-7-yl) carbamate as an intermediate**

This compound was prepared in the manner of Step C, Example 7, with 2.0 grams (8.7 mmole) of 7-amino-2-methylthio-4-chlorobenzothiazole, 40 mL pyridine, and 1.84 grams (17 mmole) of ethyl chloroformate as reagents. The yield of ethyl N-(2-methylthio-4-chlorobenzothiazol-7-yl)carbamate was 2.4 grams. The NMR spectrum was consistent with the proposed structure.

5 Step C Synthesis of 3-(2-methylthio-4-chlorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

This compound was prepared in the manner of Step E, Example 1, with 1.8 grams (5.9 mmole) of ethyl N-(2-methylthio-4-chlorobenzothiazol-7-yl)-carbamate, 1.3 grams (7.0 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate, 0.35 gram (6.5 mmole) of sodium methoxide, 25 mL of N,N-dimethylformamide, and 0.76 gram (5 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene as reagents. The yield of 3-(2-methylthio-4-chlorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 0.9 gram. The NMR spectrum was consistent with the proposed structure.

15 Step D Synthesis of 3-(2-methylthio-4-chlorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione (Compound 8)

This compound was prepared in the manner of Step F, the alternative method, Example 1, with 0.90 gram (2.3 mmole) of 3-(2-methylthio-4-chlorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 0.48 gram (3.5 mmole) of potassium carbonate, 0.50 gram (3.5 mmole) of iodomethane, and 20 mL of acetone as reagents. The yield of 3-(2-methylthio-4-chlorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 0.6 gram, m.p. 79-81 °C. The NMR spectrum was consistent with the proposed structure.

EXAMPLE 10

30 SYNTHESIS OF 3-(4-METHOXY-2,6-DICHLOROBENZOTHAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)-PYRIMIDINEDIONE (COMPOUND 18)

Step A Synthesis of 2,6-dichloro-7-nitrobenzothiazole as an intermediate

This compound was prepared in the manner of Step D, Example 6, with 8.0 grams (39 mmole) of 2,6-dichlorobenzothiazole, 45 mL of concentrated

35

sulfuric acid, and 3.8 mL (60 mmole) of concentrated nitric acid as reagents. The material was purified by column chromatography on silica gel. Elution was accomplished with 3:1 heptane:ethyl acetate. The product-containing fractions were combined and concentrated under reduced pressure, yielding
5 2,6-dichloro-7-nitrobenzothiazole. The NMR spectrum was consistent with the proposed structure.

Step B Synthesis of 7-amino-2,6-dichlorobenzothiazole as an intermediate

This compound was prepared in the manner of Step E, Example 6, with
10 9.5 grams (33 mmole) of 2,6-dichloro-7-nitrobenzothiazole, 4.5 grams (80 mmole) of iron powder, 30 mL of acetic acid, 150 mL of ethanol, and 3.3 mL (40 mmole) of concentrated hydrochloric acid as reagents. The yield of 7-amino-2,6-dichlorobenzothiazole was 6.0 grams. The NMR spectrum was consistent with the proposed structure.

15 Step C Synthesis of 7-amino-2,4,6-trichlorobenzothiazole as an intermediate

Under a nitrogen atmosphere a solution of 6.0 grams (27 mmole) of 7-amino-2,6-dichlorobenzothiazole in 75 mL of N,N-dimethylformamide was stirred and cooled in an ice bath. To this was then added 4.3 grams (32
20 mmole) of N-chlorosuccinimide. Upon completion of the addition the reaction mixture was stirred for 20 minutes and then warmed to ambient temperature and stirred for an additional 18 hours. After this time the reaction mixture was poured into 250 mL of ice-water, and about 15 grams of lithium chloride was added, after which the reaction mixture was stirred for
25 30 minutes. The resulting precipitate was collected by filtration and dried under reduced pressure, yielding a dark solid, which was purified by column chromatography on silica gel. Elution was accomplished with 4:1 hexane:ethyl acetate. The product-containing fractions were combined and concentrated under reduced pressure, yielding 4.4 grams of 7-amino-2,4,6-trichlorobenzothiazole. The NMR spectrum was consistent with the
30 proposed structure.

Step D Synthesis of (2,4,6-trichlorobenzothiazol-7-yl)isocyanate as an intermediate

Under a nitrogen atmosphere 25 mL of ethyl acetate was stirred and
35 cooled in an ice bath. To this was added via syringe 9.5 mL (18 mmole) of

1.9 M phosgene in toluene. During 25 minutes, a solution of 3.0 grams (11.8 mmole) of 7-amino-2,4,6-trichlorobenzothiazole in 75 mL of ethyl acetate was added dropwise. Upon completion of the addition the reaction mixture was warmed to ambient temperature and then gently heated at reflux for 2.5 hours. After this time the reaction mixture was concentrated under reduced pressure, yielding (2,4,6-trichlorobenzothiazol-7-yl)isocyanate, which was stored in a refrigerator.

Step E Synthesis of 3-(2,4,6-trichlorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

10 This compound was prepared in the manner of Step E, Example 1, with 2.5 grams (8.9 mmole) of (2,4,6-trichlorobenzothiazol-7-yl)isocyanate, 1.8 grams (10 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate, 0.59 gram (11 mmole) of sodium methoxide, 50 mL of N,N-dimethylformamide, and 1.5 grams (9 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene as reagents. The

15 yield of 3-(2,4,6-trichlorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione was 0.8 gram. The NMR spectrum was consistent with the proposed structure.

Step F Synthesis of 3-(4-methoxy-2,6-dichlorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione

20 (45Compound 18)

 This compound was prepared in the manner of Step F, the alternative method, Example 1, with 0.70 gram (1.8 mmole) of 3-(2,4,6-trichlorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 0.5 gram (3.6 mmole) of potassium carbonate, 0.51 gram (3.6 mmole) of iodomethane,

25 and 30 mL of acetone as reagents. The yield of 3-(4-methoxy-2,6-dichlorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 0.3 gram, m.p. 185-187 °C. The NMR spectrum indicated that the material was 3-(4-methoxy-2,6-dichlorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione rather than 3-(2,4,6-trichlorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione,

30 which was the intended compound.

EXAMPLE 11

**SYNTHESIS OF 3-(2-METHYL-4-CHLORO-6-FLUOROBENZO-
THIAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)-
PYRIMIDINEDIONE (COMPOUND 12)**

5 Step A Synthesis of 2-methyl-6-fluoro-7-nitrobenzothiazole as an intermediate

10 This compound was prepared in the manner of Step D, Example 6, with 9.3 grams (55 mmole) of 2-methyl-6-fluorobenzothiazole, 50 mL of concentrated sulfuric acid, and 4.2 mL (67 mmole) of concentrated nitric acid as reagents. The material was purified by column chromatography on silica gel. Elution was accomplished with 3:1 heptane:ethyl acetate. The product-containing fractions were combined and concentrated under reduced pressure. The yield of 2-methyl-6-fluoro-7-nitrobenzothiazole was 5.7 grams. The NMR spectrum was consistent with the proposed structure.

15 Step B Synthesis of 7-amino-2-methyl-6-fluorobenzothiazole as an intermediate

This compound was prepared in the manner of Step E, Example 6, with 7.0 grams (33 mmole) of 2-methyl-6-fluoro-7-nitrobenzothiazole, 7.0 grams (125 mmole) of iron powder, 30 mL of acetic acid, and 75 mL of ethanol as reagents. The yield of 7-amino-2-methyl-6-fluorobenzothiazole was 5.1 grams. The NMR spectrum was consistent with the proposed structure.

Step C **Synthesis of ethyl N-(2-methyl-6-fluorobenzothiazol-7-yl) carbamate as an intermediate**

25 This compound was prepared in the manner of Step C, Example 7, with 5.0 grams (27 mmole) of 7-amino-2-methyl-6-fluorobenzothiazole, 50 mL pyridine, and 5.4 grams (50 mmole) of ethyl chloroformate as reagents. This preparation differs in that the ethyl chloroformate was added to the 7-amino-2-methyl-6-fluorobenzothiazole /pyridine solution rather than the 7-amino-2-methyl-6-fluorobenzothiazole /pyridine solution being added to the ethyl
30 chloroformate. The material was purified by recrystallization from 5:1 petroleum ether and chloroform. The yield of ethyl N-(2-methyl-6-fluorobenzothiazol-7-yl)carbamate was 3.5 grams, m.p. 132-134 °C. The NMR spectrum was consistent with the proposed structure.

35 **Step D Synthesis of ethyl N-(2-methyl-4-chloro-6-fluorobenzothiazol-7-yl)carbamate as an intermediate**

5 This compound was prepared in the manner of Step D, Example 1, with 2.5 grams (9.8 mmole) of ethyl N-(2-methyl-6-fluorobenzothiazol-7-yl)carbamate, 1.6 grams (10 mmole) of N,N-dichlorourethane, 0.17 mL (2.0 mmole) of concentrated hydrochloric acid, and 35 mL of acetic acid as reagents. The yield of ethyl N-(2-methyl-4-chloro-6-fluorobenzothiazol-7-yl)carbamate was 2.2 grams, m.p. 167-169 °C. The NMR spectrum was consistent with the proposed structure.

Step E Synthesis of 3-(2-methyl-4-chloro-6-fluorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

10 This compound was prepared in the manner of Step E, Example 1, with 1.8 grams (6.2 mmole) of ethyl N-(2-methyl-4-chloro-6-fluorobenzothiazol-7-yl)carbamate, 1.25 grams (6.8 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate, 0.35 gram (6.5 mmole) of sodium methoxide, 50 mL of N,N-dimethylformamide, and 0.94 gram (6.2 mmole) of 1,8-diazabicyclo[5.4.0]-undec-7-ene as reagents. The yield of 3-(2-methyl-4-chloro-6-fluorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 1.25 grams. The NMR spectrum was consistent with the proposed structure.

Step F Synthesis of 3-(2-methyl-4-chloro-6-fluorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione
20 (Compound 12)

This compound was prepared in the manner of Step F, Example 1, with 0.55 gram (1.4 mmole) of 3-(2-methyl-4-chloro-6-fluorobenzothiazol-7-yl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 0.34 gram (2.5 mmole) of potassium carbonate, 0.35 gram (2.5 mmole) of iodomethane, and 25 mL of acetone as reagents. The yield of 3-(2-methyl-4-chloro-6-fluorobenzothiazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 0.45 gram, m.p. 83-85 °C. The NMR spectrum was consistent with the proposed structure.

30

EXAMPLE 12

SYNTHESIS OF 3-[2-(1-PIPERIDINYL)-4-CHLORO-6-FLUORO-
BENZOTHIAZOL-7-YL]-1-METHYL-6-TRIFLUOROMETHYL-2,4-
(1H,3H)PYRIMIDINEDIONE (COMPOUND 15)

Step A Synthesis of ethyl N-(2,4-dichloro-6-fluorobenzothiazol-7-yl)
35 carbamate as an intermediate

This compound was prepared in the manner of Step D, Example 1, with 3.6 grams (13.1 mmole) of ethyl N-(2-chloro-6-fluorobenzothiazol-7-yl)carbamate (prepared as in Steps A-D, Example 7), 2.2 grams (14.0 mmole) of N,N-dichlorourethane, 0.17 mL (2.0 mmole) of concentrated hydrochloric acid, and 100 mL of acetic acid as reagents. The yield of ethyl N-(2,4-dichloro-6-fluorobenzothiazol-7-yl)carbamate was 3.1 grams. The NMR spectrum was consistent with the proposed structure.

Step B Synthesis of ethyl N-[2-(1-piperidiny)-4-chloro-6-fluorobenzothiazol-7-yl]carbamate as an intermediate

Under a nitrogen atmosphere a solution of 3.0 grams (9.7 mmole) of ethyl N-(2,4-dichloro-6-fluorobenzothiazol-7-yl)carbamate in 75 mL of tetrahydrofuran was stirred, and 1.7 grams (20 mmole) of piperidine was added. Upon completion of the addition the reaction mixture was stirred for 1.5 hours at ambient temperature. The reaction mixture was then poured into a mixture of 75 mL of aqueous 10% ammonium chloride and 75 mL of an aqueous saturated sodium chloride solution. The resulting mixture was extracted with three 75 mL portions of ethyl acetate. The combined extracts were washed with two 50 mL portions of an aqueous saturated sodium chloride solution. The organic layer was dried with sodium sulfate, filtered, and the filtrate was concentrated under reduced pressure, yielding a brown solid, which was purified by column chromatography on silica gel. Elution was accomplished with 2:1 heptane:ethyl acetate. The product-containing fractions were combined and concentrated, yielding 2.6 grams of ethyl N-[2-(1-piperidiny)-4-chloro-6-fluorobenzothiazol-7-yl]carbamate. The NMR spectrum was consistent with the proposed structure.

Step C Synthesis of 3-[2-(1-piperidiny)-4-chloro-6-fluorobenzothiazol-7-yl]-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

This compound was prepared in the manner of Step E, Example 1, with 2.2 grams (6.1 mmole) of ethyl N-[2-(1-piperidiny)-4-chloro-6-fluorobenzothiazol-7-yl]carbamate, 1.3 grams (7.0 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate, 0.38 gram (7.0 mmole) of sodium methoxide, 50 mL of N,N-dimethylformamide, and 0.93 gram (6.1 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene as reagents. The yield of 3-[2-(1-piperidiny)-4-chloro-6-

fluorobenzothiazol-7-yl]-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 0.75 gram. The NMR spectrum was consistent with the proposed structure.

Step D Synthesis of 3-[2-(1-piperidiny)-4-chloro-6-fluorobenzothiazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione
(Compound 15)

5 This compound was prepared in the manner of Step F, the alternative method, Example 1, with 0.7 gram (1.6 mmole) of 3-[2-(1-piperidiny)-4-chloro-6-fluorobenzothiazol-7-yl]-6-trifluoromethyl-2,4-(1H,3H)pyrimidine-
10 dione, 0.41 gram (3.0 mmole) of potassium carbonate, 0.43 gram (3.0 mmole) of iodomethane, and 45 mL of acetone as reagents. The yield of 3-[2-(1-piperidiny)-4-chloro-6-fluorobenzothiazol-7-yl]-1-methyl-6-trifluoro-
 methyl-2,4-(1H,3H)pyrimidinedione was 0.5 gram. The NMR spectrum was consistent with the proposed structure.

15

EXAMPLE 13

SYNTHESIS OF 3-[2-(1-PIPERIDINY)-4-CHLOROBENZOTHAZOL-7-
YL]-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)-
PYRIMIDINEDIONE (COMPOUND 9)

Step A Synthesis of 2-(1-piperidiny)-4-chloro-7-nitrobenzothiazole as
20 an intermediate

 Under a nitrogen atmosphere a solution of 1.7 grams (6.8 mmole) of 2,4-dichloro-7-nitrobenzothiazole in 150 mL of diethyl ether was stirred as 1.2 grams (13.7 mmole) of piperidine was added via syringe over a ten minute period. The reaction mixture, which had spontaneously warmed, was cooled
25 with ice-water to ambient temperature, where it stirred for 40 minutes. The resulting precipitate was collected by filtration and washed with 50 mL of ethyl acetate. The filtrate was washed with aqueous 0.1 N hydrochloric acid and two 50 mL portions of water. The organic layer was dried with sodium sulfate, filtered, and concentrated under reduced pressure, yielding a bright
30 yellow-orange solid, which was combined with a material prepared by a similar route and purified by column chromatography on silica gel. Elution was accomplished with 7:1 followed by 5:1 hexane:ethyl acetate. The product-containing fractions were combined and concentrated under reduced pressure, yielding 1.3 grams of 2-(1-piperidiny)-4-chloro-7-nitro-

benzothiazole. The NMR spectrum was consistent with the proposed structure.

Step B Synthesis of 7-amino-2-(1-piperidiny)-4-chlorobenzothiazole as an intermediate

5 This compound was prepared in the manner of Step E, Example 6, with 1.2 grams (4.0 mmole) of 2-(1-piperidiny)-4-chloro-7-nitrobenzothiazole, 0.89 gram (16.0 mmole) of iron powder, 0.33 mL (4 mmole) of concentrated hydrochloric acid, 2 mL of acetic acid, and 50 mL of ethanol as reagents. The yield of 7-amino-2-(1-piperidiny)-4-chlorobenzothiazole was 1.1 grams.

10 The NMR spectrum was consistent with the proposed structure.

Step C Synthesis of ethyl N-[2-(1-piperidiny)-4-chlorobenzothiazol-7-yl]carbamate as an intermediate

15 This compound was prepared in the manner of Step C, Example 7, with 1.0 gram (3.7 mmole) of 7-amino-2-(1-piperidiny)-4-chlorobenzothiazole, 20 mL pyridine, and 0.76 gram (7.0 mmole) of ethyl chloroformate as reagents. The yield of ethyl N-[2-(1-piperidiny)-4-chlorobenzothiazol-7-yl]carbamate was 1.6 grams. The NMR spectrum was consistent with the proposed structure.

Step D Synthesis of 3-[2-(1-piperidiny)-4-chlorobenzothiazol-7-yl]-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

20 This compound was prepared in the manner of Step E, Example 1, with 1.3 grams (3.8 mmole) of ethyl N-[2-(1-piperidiny)-4-chlorobenzothiazol-7-yl]carbamate, 0.77 gram (4.2 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate, 0.22 gram (4.1 mmole) of sodium methoxide, 25 mL of N,N-dimethylformamide, and 0.5 mL (3.3 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene as reagents. The yield of 3-[2-(1-piperidiny)-4-chlorobenzothiazol-7-yl]-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 0.68 gram. The NMR spectrum was consistent with the proposed structure.

Step E Synthesis of 3-[2-(1-piperidiny)-4-chlorobenzothiazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione (Compound 9)

30 This compound was prepared in the manner of Step F, the alternative method, Example 1, with 0.66 gram (1.6 mmole) of 3-[2-(1-piperidiny)-4-chlorobenzothiazol-7-yl]-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 0.33 gram (2.4 mmole) of potassium carbonate, 0.43 gram (3.0 mmole) of iodo-

methane, and 25 mL of acetone as reagents. The yield of 3-[2-(1-piperidinyl)-4-chlorobenzothiazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione was 0.3 gram, m.p. 102-104 °C. The NMR spectrum was consistent with the proposed structure.

5

EXAMPLE 14

SYNTHESIS OF 3-(2-DIBROMOMETHYL-4-CHLORO-6-FLUORO-BENZOXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)-PYRIMIDINEDIONE (COMPOUND 56)

10 A stirred solution of 1.6 grams (4.2 mmole) of 3-(2-methyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidine-
dione, 5.0 grams (28 mmole) of N-bromosuccinimide, 0.05 gram (0.3
mmole) of 2,2'-azobis(2-methylpropionitrile), and 0.05 gram (0.2 mmole) of
benzoyl peroxide in 50 mL of carbon tetrachloride was heated to reflux,
15 where it stirred for 32 hours. After this time the reaction mixture was allowed
to cool to ambient temperature, where it stirred for about an additional 48
hours. At the conclusion of this period 50 mL of trichloromethane was
added, and the resulting precipitate was collected by filtration and washed
20 with one 20 mL portion of trichloromethane. The combined filtrate and wash
were concentrated under reduced pressure to a yellow solid, which was
subjected to column chromatography on silica gel. Elution was with 5:1
followed by 2:1 heptane and ethyl acetate. The product-containing fractions
were combined and concentrated under reduced pressure, yielding 0.15
25 gram of 3-(2-dibromomethyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-
trifluoromethyl-2,4-(1H,3H)pyrimidinedione. The NMR spectrum was
consistent with the proposed structure. This 0.15 gram of product was
combined with 0.39 gram of product prepared by a similar route, yielding a
total of 0.54 gram of 3-(2-dibromomethyl-4-chloro-6-fluorobenzoxazol-7-yl)-
1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, m.p. 144-146°C.

30

EXAMPLE 15

SYNTHESIS OF 3-[2-(1-HYDROXY-1-METHYLETHYL)-4-CHLORO-6-FLUOROBENZOXAZOL-7-YL]-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE (COMPOUND 58)

Step A Synthesis of N-(3-chloro-4-fluorophenyl)-2-methoxy-2-methylpropionamide as an intermediate

A stirred solution of 6.1 grams (42 mmole) of 3-chloro-4-fluoroaniline, 5.0 grams (42 mmole) of 2-methoxy-2-methylpropanoic acid, and 10.5 grams (55 mmole) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride in 120 mL of methylene chloride was cooled to about 0 °C, and 7.4 grams (55 mmole) of 1-hydroxybenzotriazole was added. Upon completion of the addition the reaction was slowly warmed to ambient temperature, where it stirred for about 18 hours. After this time the reaction mixture was poured into water, and the resulting mixture was filtered through diatomaceous earth to remove any salts that had precipitated. The organic layer was separated from the aqueous layer, washed with an aqueous saturated sodium chloride solution, dried with sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to a brown oil, which was passed through a silica gel scrub plug, yielding 5.0 grams of N-(3-chloro-4-fluorophenyl)-2-methoxy-2-methylpropionamide. The NMR spectrum was consistent with the proposed structure.

Step B Synthesis of [2-(1-methoxy-1-methylethyl)-6-fluorobenzoxazol-7-yl]carboxylic acid as an intermediate

This compound was prepared in the manner of Step B, Example 1, with 4.9 grams (20 mmole) of N-(3-chloro-4-fluorophenyl)-2-methoxy-2-methylpropionamide, 120 mL of tetrahydrofuran, 17.6 mL (44 mmole) of 2.5 M n-butyllithium in hexanes, and 3.0 grams of solid carbon dioxide as reagents. The yield of [2-(1-methoxy-1-methylethyl)-6-fluorobenzoxazol-7-yl]carboxylic acid was 3.5 grams. The NMR spectrum was consistent with the proposed structure.

An additional 2.1 grams of [2-(1-methoxy-1-methylethyl)-6-fluorobenzoxazol-7-yl]carboxylic acid of was prepared in the manner of Step B, Example 1, with 4.2 grams (17 mmole) of N-(3-chloro-4-fluorophenyl)-2-methoxy-2-methylpropionamide, 120 mL of tetrahydrofuran, 16 mL (40 mmole) of 2.5 M n-butyllithium in hexanes, and 3.0 grams of solid carbon dioxide as reagents. The NMR spectrum was consistent with the proposed structure.

Step C Synthesis of [2-(1-methoxy-1-methylethyl)-6-fluorobenzoxazol-7-yl]carbonyl azide as an intermediate

This compound was prepared in the manner of Step C, Example 1, with 5.6 grams (22 mmole) of [2-(1-methoxy-1-methylethyl)-6-fluorobenzoxazol-7-yl]carboxylic acid, 30 mL of tetrahydrofuran, 2.3 grams (22.7 mmole) of 4-methylmorpholine, 2.2 mL (27 mmole) of ethyl chloroformate, 3.0 grams (45 mmole) of sodium azide, and 3 mL of water as reagents. The yield of [2-(1-methoxy-1-methylethyl)-6-fluorobenzoxazol-7-yl]carbonyl azide was 3.7 grams. The NMR spectrum was consistent with the proposed structure.

Step D Synthesis of ethyl N-[2-(1-methoxy-1-methylethyl)-6-fluorobenzoxazol-7-yl]carbamate as an intermediate

Under a nitrogen atmosphere a stirred solution of 3.7 grams (13 mmole) of [2-(1-methoxy-1-methylethyl)-6-fluorobenzoxazol-7-yl]carbonyl azide in 25 mL of ethanol was heated to reflux where it stirred for about 18 hours. After this time the reaction mixture was cooled to ambient temperature, and the ethanol was removed, yielding a yellow oil, which was subjected to column chromatography on silica gel. Elution was with 4:1 heptane and ethyl acetate. The product-containing fractions were combined and concentrated under reduce pressure, yielding 2.5 grams of ethyl N-[2-(1-methoxy-1-methylethyl)-6-fluorobenzoxazol-7-yl]carbamate. The NMR spectrum was consistent with the proposed structure.

Step E Synthesis of ethyl N-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]carbamate as an intermediate

This compound was prepared in the manner of Step D, Example 1, with 2.5 grams (8.8 mmole) of ethyl N-[2-(1-methoxy-1-methylethyl)-6-fluorobenzoxazol-7-yl]carbamate, 30 mL of acetic acid, about 0.5 mL (6 mmole) of concentrated hydrochloric acid, and 1.4 grams (8.8 mmole) of N,N-dichloro-urethane as reagents. The yield of ethyl N-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]carbamate was 2.2 grams, m.p. 95-97 °C. The NMR spectrum was consistent with the proposed structure.

Step F Synthesis of 3-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

This compound was prepared in the manner of Step E, Example 1, with 0.8 gram (2.4 mmole) of ethyl N-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]carbamate, 0.15 gram (2.8 mmole) of sodium methoxide, 0.44 gram (2.4 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate, 30 mL of

5 N,N-dimethylformamide, and 0.37 gram (2.4 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene as reagents. The yield of 3-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione was 0.3 gram. The NMR spectrum was consistent with the proposed structure.

10 An additional 0.6 gram of 3-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was prepared in the manner of Step E, Example 1, with 0.8 gram (2.4 mmole) of ethyl N-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]carbamate, 0.15 gram (2.8 mmole) of sodium methoxide, 0.44 gram (2.4 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate, 30 mL of N,N-dimethylformamide, and 0.37 gram (2.4 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene as reagents. The NMR spectrum was consistent with the proposed structure.

15 **Step G** Synthesis of 3-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione as an intermediate

20 This compound was prepared in the manner of Step F, Example 1, with 0.6 gram (1.4 mmole) of 3-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 0.39 gram (2.8 mmole) of potassium carbonate, 0.4 gram (2.8 mmole) of methyl iodide, and 15 mL of acetone as reagents. The yield of 3-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 0.6 gram. The NMR spectrum was consistent with the proposed structure.

25 **Step H** Synthesis of 3-[2-(1-hydroxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione (Compound 58)

30 Under a nitrogen atmosphere a stirred solution of 3.8 mL (3.8 mmole) of 1 M boron tribromide (in methylene chloride) in 50 mL of methylene chloride was cooled to about -40 °C, and a solution of 0.6 gram (2.6 mmole) of 3-[2-(1-methoxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione in 50 mL of methylene chloride was added dropwise at such a rate as to maintain the reaction mixture temperature below -30 °C. Upon completion of the addition the reaction

35

mixture was allowed to warm to ambient temperature, where it stirred for about 18 hours. At the conclusion of this period the reaction mixture was poured into water and thoroughly extracted with methylene chloride. The combined extracts were washed with an aqueous saturated sodium chloride solution, dried with sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to a brown oil, which was subjected to column chromatography on silica gel. Elution was with 1:4 ethyl acetate to heptane. The product-containing fractions were combined and concentrated under reduced pressure, yielding 0.3 gram of 3-[2-(1-hydroxy-1-methylethyl)-4-chloro-6-fluorobenzoxazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione. The NMR spectrum was consistent with the proposed structure.

EXAMPLE 16

SYNTHESIS OF 3-(2-THIO-4-CHLORO-6-FLUOROBENZOXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE (COMPOUND 53)

Step A Synthesis of ethyl N-(4-chloro-2,6-difluorophenyl)carbamate as an intermediate

This compound was prepared in the manner of Step A, Example 6, with 26.9 grams (165 mmole) of 4-chloro-2,6-difluoroaniline and 17.4 mL (182 mmole) of ethyl chloroformate in 90 mL of pyridine as reagents. The yield of ethyl N-(4-chloro-2,6-difluorophenyl)carbamate was 23.6 grams. The NMR spectrum was consistent with the proposed structure.

Step B Synthesis of ethyl N-(4-chloro-2,6-difluoro-3-nitrophenyl)-carbamate as an intermediate

This compound was prepared in the manner of Step D, Example 6, with 23.6 grams (109 mmole) of ethyl N-(4-chloro-2,6-difluorophenyl)carbamate and 7.7 mL (123 mmole) of 70% nitric acid in 125 mL of sulfuric acid as reagents. The yield of ethyl N-(4-chloro-2,6-difluoro-3-nitrophenyl)carbamate is 30.6 grams. The NMR spectrum was consistent with the proposed structure.

Step C Synthesis of ethyl N-(4-chloro-6-fluoro-2-methoxy-3-nitrophenyl)carbamate as an intermediate

Under a nitrogen atmosphere a solution of 30.6 grams (109 mmole) of ethyl N-(4-chloro-2,6-difluoro-3-nitrophenyl)carbamate and 18 mL (449 mmole) of methanol in 175 mL of dioxane was stirred, and 218 mL (218 mmole) of 1 M sodium trimethylsilanoate (in tetrahydrofuran) was added dropwise during a 45 minute period. Upon completion of the addition the reaction mixture was heated to 65 °C, where it stirred for three hours. At the conclusion of this period the reaction mixture was allowed to cool to ambient temperature, where it stirred for about 18 hours. The reaction mixture was concentrated under reduced pressure to a residue, which was taken up in cold 3 N hydrochloric acid. The resulting solid was collected by filtration, washed with petroleum ether, and heat dried under vacuum, yielding 21.3 grams of ethyl N-(4-chloro-6-fluoro-2-methoxy-3-nitrophenyl)carbamate. The NMR spectrum was consistent with the proposed structure.

Step D Synthesis of ethyl N-(3-amino-4-chloro-6-fluoro-2-methoxyphenyl)carbamate as an intermediate

This compound was prepared in the manner of Step E, Example 6, with 21.3 grams (72 mmole) of ethyl N-(4-chloro-6-fluoro-2-methoxy-3-nitrophenyl)carbamate, 18.3 grams (328 mmole) of iron powder, 50 mL of acetic acid, 250 mL of ethanol, and 3 mL (36 mmole) of hydrochloric acid as reagents. This preparation differed in that the hydrochloric acid was added. The yield of ethyl N-(3-amino-4-chloro-6-fluoro-2-methoxyphenyl)carbamate was 15.0 grams. The NMR spectrum was consistent with the proposed structure.

Step E Synthesis of 3-(3-amino-4-chloro-6-fluoro-2-methoxyphenyl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

This compound was prepared in the manner of Step E, Example 1, with 15.0 grams (57 mmole) of ethyl N-(3-amino-4-chloro-6-fluoro-2-methoxyphenyl)carbamate, 3.6 grams (66 mmole) of sodium methoxide, and 10.5 grams (57 mmole) of ethyl 3-amino-4,4,4-trifluorocrotonate in 100 mL of N,N-dimethylformamide as reagents. This preparation differed in that no 1,8-diazabicyclo[5.4.0]undec-7-ene was added. The yield of 3-(3-amino-4-chloro-6-fluoro-2-methoxyphenyl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidine-dione was 10.0 grams. The NMR spectrum was consistent with the proposed structure.

Step F Synthesis of 3-(3-amino-4-chloro-6-fluoro-2-methoxyphenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

5 This compound was prepared in the manner of Step F, Example 1, with 10.0 grams (28 mmole) of 3-(3-amino-4-chloro-6-fluoro-2-methoxyphenyl)-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 5.7 grams (41 mmole) of potassium carbonate, 26 mL (41 mmole) of methyl iodide, and 70 mL of acetone as reagents. The yield of 3-(3-amino-4-chloro-6-fluoro-2-methoxyphenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 7.3
10 grams. The NMR spectrum was consistent with the proposed structure.

Step G Synthesis of 3-(3-amino-4-chloro-6-fluoro-2-hydroxy)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione as an intermediate

15 This compound was prepared in the manner of Step H, Example 15, with 7.3 grams (20 mmole) of 3-(3-amino-4-chloro-6-fluoro-2-methoxyphenyl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione and 80 mL (80 mmole) of 1 M boron tribromide (in methylene chloride) in 120 mL of methylene chloride as reagents. The yield of 3-(3-amino-4-chloro-6-fluoro-2-hydroxy)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 5.4
20 grams. The NMR spectrum was consistent with the proposed structure.

Step H Synthesis of 3-(2-thio-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione
(Compound 53)

25 Under a nitrogen atmosphere a stirred solution of 5.4 grams (15 mmole) of 3-(3-amino-4-chloro-6-fluoro-2-hydroxy)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione and 4.9 grams (30 mmole) of potassium ethyl xanthate in 25 mL of ethanol was heated at reflux for four hours. After this time the reaction mixture was cooled to ambient temperature and concentrated under reduced pressure to a residue, which was taken up in water and
30 acidified to pH 5 with acetic acid. The resulting mixture was thoroughly extracted with methylene chloride. The combined extracts were washed with an aqueous saturated sodium chloride solution, dried with sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to a brown oil, which was subjected to column chromatography on silica gel.
35 Elution was with 1:3 followed by 1:1 ethyl acetate to heptane. The product-

containing fractions were combined and concentrated under reduced pressure, yielding 1.0 gram of crude product. This crude product was purified by column chromatography on silica gel. Elution was with 1:9 methanol and methylene chloride. The product-containing fractions were
5 combined and concentrated under reduced pressure, yielding 0.3 gram of 3-(2-thio-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione. The NMR spectrum was consistent with the proposed structure.

10

EXAMPLE 17

SYNTHESIS OF 3-(2-METHYLTHIO-4-CHLORO-6-FLUOROBENZOXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)-PYRIMIDINEDIONE (COMPOUND 51)

This compound was prepared in the manner of Step F, Example 1, with
15 4.0 grams (10 mmole) of 3-(2-thio-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 1.5 grams (11 mmole) of potassium carbonate, 1.7 grams (12 mmole) of methyl iodide, and 25 mL of acetone as reagents. The yield of 3-(2-methylthio-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione was 0.4
20 gram. The NMR spectrum was consistent with the proposed structure.

EXAMPLE 18

SYNTHESIS OF 3-(2-AMINO-4-CHLORO-6-FLUOROBENZOXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE (COMPOUND 70)

25

A stirred solution of 1.3 grams (3.5 mmole) of 3-(3-amino-4-chloro-6-fluoro-2-hydroxy)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione in 20 mL of methanol was cooled to about 0 °C in an ice bath and 0.4 gram (3.5 mmole) of cyanogen bromide was added dropwise during a ten minute
30 period. At the conclusion of this period the reaction mixture was warmed to ambient temperature, where it stirred for about 18 hours. The reaction mixture was poured into an aqueous saturated sodium bicarbonate solution. The resulting mixture was thoroughly extracted with portions of ethyl acetate. The combined extracts were washed with an aqueous saturated sodium
35 chloride solution, dried with sodium sulfate, and filtered. The filtrate was

concentrated under reduced pressure to a dark oil, which was subjected to column chromatography on silica gel. Elution was with 1:50 methanol and methylene chloride. The product-containing fractions were combined and concentrated under reduced pressure to yield the crude product, which was
5 purified by recrystallization from methylene chloride. The yield of 3-(2-amino-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 0.5 grams, m.p. 144-150 °C. The NMR spectrum was consistent with the proposed structure.

10

EXAMPLE 19

SYNTHESIS OF 3-(2-ETHOXYACETYL-4-CHLORO-6-FLUOROBENZ-
OXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)-
PYRIMIDINEDIONE (COMPOUND 93)

Step A Synthesis of 3-[5-(ethylmalonylamino)-6-(ethylmalonyloxy)-4-
15 chloro-2-fluorophenyl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-
pyrimidinedione as an intermediate

Under a nitrogen atmosphere a stirred solution of 1.2 grams (8.4 mmole) of phosphorus pentoxide and 3.2 grams (20 mmole) of hexamethyl-
disiloxane was heated to 80 °C, and a solution of 1.5 grams (4.2 mmole) of
20 3-(3-amino-4-chloro-6-fluoro-2-hydroxy)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione and 1.0 gram (6 mmole) of the potassium salt of ethyl malonate in 50 mL of dioxane was added. Upon completion of the addition the reaction mixture was heated to reflux, where it stirred for 16
25 hours. After this time the reaction mixture was cooled to ambient temperature and poured into 100 mL of water. The resulting mixture was extracted with three 50 mL portions of ethyl acetate. The extracts were combined and washed with two 50 mL portions of an aqueous saturated sodium chloride solution. The combined extracts and washes were dried with magnesium sulfate and filtered. The filtrate was concentrated under reduced pressure,
30 yielding 2.6 grams of a yellow oil. The NMR spectrum indicated that the yellow oil was 3-[5-(ethylmalonylamino)-6-(ethylmalonyloxy)-4-chloro-2-fluorophenyl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione rather than 3-(2-ethoxyacetyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione.

Step B Synthesis of 3-(2-ethoxyacetyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione (Compound 93)

This compound was prepared in the manner of Step G, Example 6, with
5 2.6 grams (4.4 mmole) of 3-[5-(ethylmalonylamino)-6-(ethylmalonyloxy)-4-chloro-2-fluorophenyl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidine-
dione, 1.2 grams (8.4 mmole) of phosphorus pentoxide, and 3.2 grams (20
mmole) of hexamethyldisiloxane in 30 mL of toluene as reagents. This
preparation differed in that toluene was used rather than acetic acid. The
10 yield of 3-(2-ethoxyacetyl-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione was 0.5 gram, m.p. 56-58 °C.
The NMR spectrum was consistent with the proposed structure.

EXAMPLE 20

15 **SYNTHESIS OF 3-(2-ACETYLAMINO-4-CHLORO-6-FLUOROBENZ-
OXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)-
PYRIMIDINEDIONE (COMPOUND 115)**

A solution of 3.8 grams (10 mmole) of 3-(2-amino-4-chloro-6-fluoro-
benzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione and
20 1.0 gram (15 mmole) of acetyl chloride in 20 mL of pyridine is stirred at
ambient temperature for about 18 hours. The reaction mixture is concentra-
ted under reduced pressure, yielding 3-(2-acetylamino-4-chloro-6-fluoro-
benzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione.

25 **EXAMPLE 21**

**SYNTHESIS OF 3-[2-(N-ACETYL-N-METHYLSULFONYLAMINO)-4-
CHLORO-6-FLUOROBENZOXAZOL-7-YL]-1-METHYL-6-
TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE
(COMPOUND 116)**

30 A stirred suspension of 2.0 grams (51 mmole) of 60% sodium hydride
(in mineral oil) in 25 mL of tetrahydrofuran is cooled in an ice bath. To this is
added a solution of 10.2 grams (25.5 mmole) of 3-(2-acetylamino-4-chloro-6-
fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidine-
dione in 25 mL of tetrahydrofuran. Upon completion of the addition 2.9
35 grams (25.3 mmole) of methanesulfonyl chloride is added. The reaction

mixture is heated to reflux, where it stirs for one hour and then concentrated under reduced pressure, yielding 3-[2-(N-acetyl-N-methylsulfonylamino)-4-chloro-6-fluorobenzoxazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinedione.

5

EXAMPLE 22

SYNTHESIS OF 3-(2-METHYLSULFONYLAMINO-4-CHLORO-6-
FLUOROBENZOXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-
(1H,3H)PYRIMIDINEDIONE (COMPOUND 71)

10 A stirred solution of 4.5 grams (10 mmole) of 3-[2-(N-acetyl-N-methylsulfonylamino)-4-chloro-6-fluorobenzoxazol-7-yl]-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione in 25 mL of 85% hydrazine is heated to 60 °C, where it stirs for about 18 hours. After this time the reaction mixture is concentrated under reduced pressure, yielding 3-(2-methylsulfonylamino-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)-
15 pyrimidinedione.

The 3-(2-methylsulfonylamino-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione can also be prepared in the manner stated above with 3 N hydrochloric acid rather than hydrazine
20 and by heating at reflux rather than 60 °C.

EXAMPLE 23

SYNTHESIS OF 3-[2-(N-METHYL-N-METHYLSULFONYLAMINO)-4-
CHLORO-6-FLUOROBENZOXAZOL-7-YL]-1-METHYL-6-
25 TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE
(COMPOUND 76)

This compound is prepared in the manner of Step F, Example 1, with 4.3 grams (10 mmole) of 3-(2-methylsulfonylamino-4-chloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, 1.5
30 grams (11 mmole) of potassium carbonate, 1.7 grams (12 mmole) of methyl iodide, and 25 mL of acetone as reagents.

EXAMPLE 24

SYNTHESIS OF 3-(2,4-DICHLORO-6-FLUOROBENZOXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE
(COMPOUND 106)

- 5 Step A Synthesis of 3-(2-oxo-4-chloro-6-fluoro-2,3-dihydrobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidine dione as an intermediate

Under a nitrogen atmosphere a stirred solution of 0.7 gram (4.3 mmole) of 1,1'-carbonyldiimidazole and 1.0 gram (2.8 mmole) of 3-(3-amino-4-chloro-6-fluoro-2-hydroxy)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidine-
10 dione in 30 mL of tetrahydrofuran was heated to reflux, where it stirred for 1.5 hours, then cooled to ambient temperature, where it stirred for about 18 hours. After this time the reaction mixture was poured into 1N aqueous hydrochloric acid. The resulting mixture was thoroughly extracted with
15 portions of ethyl acetate. The combined extracts were washed with portions of an aqueous saturated sodium chloride solution, dried with magnesium sulfate, and filtered. The filtrate was concentrated under reduced pressure, yielding 1.0 gram of 3-(2-oxo-4-chloro-6-fluoro-2,3-dihydrobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione, m.p. 67-71°C. The
20 NMR spectrum was consistent with the proposed structure.

- Step B Synthesis of 3-(2,4-dichloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione (Compound 106)


A stirring solution of 3.8 grams (10 mmole) of 3-(2-oxo-4-chloro-6-fluoro-2,3-dihydrobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimi-
25 dinedione and 7.7 grams (50 mmole) of phosphorus oxychloride is cooled in an ice bath. To this 5.0 grams (49.4 mmole) of triethylamine is added via syringe during a ten minute period. At the conclusion of this period the reaction mixture is heated to reflux where it stirs for four hours. After this period the reaction mixture is cooled to ambient temperature and poured into
30 ice water. The resulting mixture is extracted with portions of diethyl ether. The extracts are combined, washed with an aqueous saturated sodium bicarbonate solution, dried with sodium sulfate, and filtered. The filtrate is concentrated under reduced pressure, yielding 3-(2,4-dichloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione.

EXAMPLE 25

SYNTHESIS OF 3-(4-CHLORO-6-FLUORO-2-PHENOXYBENZOXAZOL-7-YL)-1-METHYL-6-TRIFLUOROMETHYL-2,4-(1H,3H)PYRIMIDINEDIONE
(COMPOUND 118)

- 5 Under a nitrogen atmosphere, a stirred solution of 2.0 grams (51 mmole) of 60% sodium hydride (in mineral oil) in 25 mL of N,N-dimethylformamide is cooled in an ice bath, and a solution of 5.3 grams (56.3 mmoles) of phenol in 25 mL of N,N-dimethylformamide is added dropwise during a five minute period. To this is added a solution of 20.3 grams (51 mmole) of 3-(2,4-dichloro-6-fluorobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione (as prepared in Example 24) in 25 mL of N,N-dimethylformamide. Upon completion of the addition the reaction mixture is allowed to warm to ambient temperature, where it stirs for about 18 hours. After this time the reaction mixture is concentrated under reduced pressure, yielding 3-(4-chloro-6-fluoro-2-phenoxybenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione.

Novel intermediates used in the preparation of the compounds described above include compounds of the formula:



20

in which: P is nitro, amino, or azido; W is hydrogen, alkoxy, or hydroxy; R¹ is hydrogen, lower alkyl, or amino; X is oxygen; Y is hydrogen, hydrogen, halogen, alkoxy, or nitro; Z is halogen; and P and W together may be -N-C(O)-O- to complete a benzoxazole ring.

25

HERBICIDAL ACTIVITY

The 3-(substituted benzoxazol-7-yl) and 3-(substituted benzothiazol-7-yl)-1-substituted-6-trifluoromethyl-2,4-(1H,3H)pyrimidinediones of the present invention were tested for pre- and postemergence herbicidal activity on a

variety of crops and weeds. The test plants included soybean (Glycine max var. Winchester), field corn (Zea mays var. Pioneer 3732), wheat (Triticum aestivum var. Lew), morningglory (Ipomea lacunosa or Ipomea hederacea), velvetleaf (Abutilon theophrasti), green foxtail (Setaria viridis), Johnsongrass (Sorghum halepense), blackgrass (Alopecurus myosuroides), common chickweed (Stellaria media), and common cocklebur (Xanthium strumarium L.).

For preemergence testing, two disposable fiber flats (8 cm x 15 cm x 25 cm) for each rate of application of each candidate herbicide were filled to an approximate depth of 6.5 cm with steam-sterilized sandy loam soil. The soil was leveled and impressed with a template to provide five evenly spaced furrows 13 cm long and 0.5 cm deep in each flat. Seeds of soybean, wheat, corn, green foxtail, and johnsongrass were planted in the furrows of the first flat, and seeds of velvetleaf, morningglory, common chickweed, cocklebur, and blackgrass were planted in the furrows of the second flat. The five-row template was employed to firmly press the seeds into place. A topping soil of equal portions of sand and sandy loam soil was placed uniformly on top of each flat to a depth of approximately 0.5 cm. Flats for postemergence testing were prepared in the same manner except that they were planted 9-14 days prior to the preemergence flats and were placed in a greenhouse and watered, thus allowing the seeds to germinate and the foliage to develop.

In both pre- and postemergence tests, a stock solution of the candidate herbicide was prepared by dissolving 0.27g of the compound in 20 mL of water/acetone (50/50) containing 0.5% v/v sorbitan monolaurate. For an application rate of 3 g/ha of herbicide a 0.10 mL portion of the stock solution was diluted with water/acetone (50/50) to 45.01 mL.

The preemergence flats were initially subjected to a light water spray. The four flats were placed two by two along a conveyor belt (i.e., the two preemergence followed by the two postemergence flats). The conveyor belt passed under a spray nozzle mounted about ten inches above the postemergent foliage. The preemergent flats were elevated on the belt so that the soil surface was at the same level below the spray nozzle as the foliage canopy of the postemergent plants. The spray of herbicidal solution was commenced, and once the spray stabilized, the flats were passed under the spray at a speed to receive a coverage equivalent of 1000L/ha, affording

the rate of 3 g/ha. The preemergence flats were watered immediately thereafter, placed in the greenhouse, and watered regularly at the soil surface. The postemergence flats were immediately placed in the greenhouse and not watered until 24 hours after treatment with the test solution.

- 5 Thereafter they were regularly watered at ground level. After 12-17 days the plants were examined and the phytotoxicity data were recorded.

Herbicidal activity data are given for various compounds of the present invention in Table 3 (preemergence) and Table 4 (postemergence). The test compounds are identified by numbers that correspond to those in Table 1.

- 10 Phytotoxicity data are taken as percent control. Percent control is determined by a method similar to the 0 to 100 rating system disclosed in "Research Methods in Weed Science," 2nd ed., B. Truelove, Ed.; Southern Weed Science Society; Auburn University, Auburn, Alabama, 1977. The rating system is as follows:

Herbicide Rating System

<u>Rating Percent Control</u>	<u>Description of Main Categories</u>	<u>Crop Description</u>	<u>Weed Description</u>
0	No effect	No crop reduction or injury	No weed control
10		Slight discoloration or stunting	Very poor weed control
20	Slight effect	Some discoloration, stunting or stand loss	Poor weed control
30		Crop injury more pronounced but not lasting	Poor to deficient weed control
40		Moderate injury, crop usually recovers	Deficient weed control
50	Moderate effect	Crop injury more lasting, recovery doubtful	Deficient to moderate weed control
60		Lasting crop injury, no recovery	Moderate weed control
70		Heavy injury and stand loss	Control somewhat less than satisfactory
80	Severe	Crop nearly destroyed, a few survivors	Satisfactory to good weed control
90		Only occasional live plants left	Very good to excellent control
100	Complete effect	Complete crop destruction	Complete weed destruction

For herbicidal application, the 3-(substituted benzoxazol-7-yl) and 3-(substituted benzothiazol-7-yl)-1-substituted-6-trifluoromethyl-2,4-(1H,3H)-pyrimidinediones are formulated into herbicidal compositions by admixture in herbicidally effective amounts with adjuvants and carriers normally employed
5 in the art for facilitating the dispersion of active ingredients for the particular utility desired, recognizing the fact that the formulation and mode of application of a toxicant may affect the activity of the material in a given application. Thus, for agricultural use the present herbicidal compounds may be formulated as granules of relatively large particle size, as water-
10 soluble or water-dispersible granules, as powdery dusts, as wettable powders, as emulsifiable concentrates, as solutions, or as any of several other known types of formulations, depending on the desired mode of application.

These herbicidal compositions may be applied either as water-diluted
15 sprays, or dusts, or granules to the areas in which suppression of vegetation is desired. These formulations may contain as little as 0.1%, 0.2% or 0.5% to as much as 95% or more by weight of active ingredient.

Dusts are free flowing admixtures of the active ingredient with finely divided solids such as talc, natural clays, kieselguhr, flours such as walnut
20 shell and cottonseed flours, and other organic and inorganic solids which act as dispersants and carriers for the toxicant; these finely divided solids have an average particle size of less than about 50 microns. A typical dust formulation useful herein is one containing 1.0 part or less of the herbicidal compound and 99.0 parts of talc.

25 Wettable powders, also useful formulations for both pre- and post-emergence herbicides, are in the form of finely divided particles which disperse readily in water or other dispersants. The wettable powder is ultimately applied to the soil either as a dry dust or as an emulsion in water or other liquid. Typical carriers for wettable powders include Fuller's earth,
30 kaolin clays, silicas, and other highly absorbent, readily wettable inorganic diluents. Wettable powders normally are prepared to contain about 5-80% of active ingredient, depending on the absorbency of the carrier, and usually also contain a small amount of a wetting, dispersing or emulsifying agent to facilitate dispersion. For example, a useful wettable powder formulation
35 contains 80.8 parts of the herbicidal compound, 17.9 parts of Palmetto clay,

1.0 part of sodium lignosulfonate, and 0.3 part of sulfonated aliphatic polyester as wetting agents. Frequently, additional wetting agent and/or oil will be added to the tank mix for postemergence application to facilitate dispersion on the foliage and absorption by the plant.

- 5 Other useful formulations for herbicidal applications are emulsifiable concentrates (ECs) which are homogeneous liquid compositions dispersible in water or other dispersant, and may consist entirely of the herbicidal compound and a liquid or solid emulsifying agent, or may also contain a liquid carrier, such as xylene, heavy aromatic naphthas, isophorone, or other
- 10 non-volatile organic solvent. For herbicidal application these concentrates are dispersed in water or other liquid carrier, and normally applied as a spray to the area to be treated. The percentage by weight of the essential active ingredient may vary according to the manner in which the composition is to be applied, but in general comprises 0.5 to 95% of active ingredient by
- 15 weight of the herbicidal composition.

Flowable formulations are similar to ECs except that the active ingredient is suspended in a liquid carrier, generally water. Flowables, like ECs, may include a small amount of a surfactant, and contain active ingredient in the range of 0.5 to 95%, frequently from 10 to 50%, by weight of

20 the composition. For application, flowables may be diluted in water or other liquid vehicle, and are normally applied as a spray to the area to be treated.

Typical wetting, dispersing or emulsifying agents used in agricultural formulations include, but are not limited to, the alkyl and alkylaryl sulfonates and sulfates and their sodium salts; alkylaryl polyether alcohols; sulfated

25 higher alcohols; polyethylene oxides; sulfonated animal and vegetable oils; sulfonated petroleum oils; fatty acid esters of polyhydric alcohols and the ethylene oxide addition products of such esters; and the addition product of long-chain mercaptans and ethylene oxide. Many other types of useful surface-active agents are available in commerce. The surface-active agent,

30 when used, normally comprises from 1 to 15% by weight of the composition.

Other useful formulations include suspensions of the active ingredient in a relatively non-volatile solvent such as water, corn oil, kerosene, propylene glycol, or other suitable solvents.

Still other useful formulations for herbicidal applications include simple

35 solutions of the active ingredient in a solvent in which it is completely soluble

at the desired concentration, such as acetone, alkylated naphthalenes, xylene, or other organic solvents. Granular formulations, wherein the toxicant is carried on relatively coarse particles, are of particular utility for aerial distribution or for penetration of cover crop canopy. Pressurized
5 sprays, typically aerosols wherein the active ingredient is dispersed in finely divided form by a propellant, such as carbon dioxide, propane or butane, may also be used. Water-soluble or water-dispersible granules are also useful formulations for herbicidal application of the present compounds. Such granular formulations are free-flowing, non-dusty, and readily water-
10 soluble or water-miscible. The soluble or dispersible granular formulations described in U.S. patent No. 3,920,442 are useful herein with the present herbicidal compounds. In use by the farmer on the field, the granular formulations, emulsifiable concentrates, flowable concentrates, solutions, etc., may be diluted with water to give a concentration of active ingredient in
15 the range of say 0.1% or 0.2% to 1.5% or 2%.


The 3-(substituted benzoxazol-7-yl) and 3-(substituted benzothiazol-7-yl)-1-substituted-6-trifluoromethyl-2,4-(1H,3H)pyrimidinediones of this invention may be formulated and/or applied with insecticides, fungicides, nematocides, plant growth regulators, fertilizers, or other agricultural
20 chemicals and may be used as effective soil sterilants as well as selective herbicides in agriculture. In applying an active compound of this invention, whether formulated alone or with other agricultural chemicals, an effective amount and concentration of the active compound is of course employed; the amount may be as low as, for example, about 4 to 300 g/ha to,
25 preferably, about 10 to 30 g/ha. For field use, where there are losses of herbicide, higher application rates (for example, four times the rates mentioned above) may be employed.

The 3-(substituted benzoxazol-7-yl) and 3-(substituted benzothiazol-7-yl)-1-substituted-6-trifluoromethyl-2,4-(1H,3H)pyrimidinediones of this
30 invention may be used in combination with other herbicides, for example they may be mixed with, say, an equal or larger amount of a known herbicide such as aryloxyalkanoic acid herbicides such as (2,4-dichlorophenoxy)acetic acid (2,4-D), (4-chloro-2-methylphenoxy)acetic acid (MCPA), (+/-)-2-(4-chloro-2-methylphenoxy)propanoic acid (MCPA); urea herbicides, such as
35 N,N-dimethyl-N'-[4-(1-methylethyl)phenyl]urea (isoproturon); imidazolinone

herbicides, such as 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid (imazapyr), a reaction product comprising (+/-)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-4-methylbenzoic acid and (+/-)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-methylbenzoic acid (imazamethabenz), (+/-)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid (imazethapyr), and (+/-)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid (imazaquin); diphenyl ether herbicides, such as 5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoic acid (acifluorfen), methyl 5-(2,4-dichlorophenoxy)-2-nitrobenzoate (bifenox), and 5-[2-chloro-4-(trifluoromethyl)phenoxy]-N-(methylsulfonyl)-2-nitrobenzamide (fomasafen); hydroxybenzonitrile herbicides, such as 4-hydroxy-3,5-diiodobenzonitrile (ioxynil), and 3,5-dibromo-4-hydroxybenzonitrile (bromoxynil); sulfonylurea herbicides, such as 2-[[(4-chloro-6-methoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]benzoic acid (chlorimuron), 2-chloro-N-[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]benzenesulfonamide (chlorsulfuron), 2-[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]methyl]benzoic acid (bensulfuron), 2-[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-1-methyl-1H-pyrazol-4-carboxylic acid (pyrazosulfuron), 3-[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylic acid (thifensulfuron), and 2-(2-chloroethoxy)-N-[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]benzenesulfonamide (triasulfuron); 2-(4-aryloxyphenoxy)alkanoic acid herbicides, such as (+/-)-2-[4-[(6-chloro-2-benzoxazolyl)oxy]phenoxy]propanoic acid (fenoxaprop), (+/-)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoic acid (fluazifop), (+/-)-2-[4-(6-chloro-2-quinoxalinyloxy)phenoxy]propanoic acid (quizalofop), and (+/-)-2-[(2,4-dichlorophenoxy)phenoxy]propanoic acid (diclofop); benzothiadiazinone herbicides, such as 3-(1-methylethyl)-1H-2,1,3-benzothiadiazin-4(3H)-one 2,2-dioxide (bentazone); 2-chloroacetanilide herbicides, such as N-(butoxymethyl)-2-chloro-2',6'-diethylacetanilide (butachlor); arenecarboxylic acid herbicides, such as 3,6-dichloro-2-methoxybenzoic acid (dicamba); and pyridyloxyacetic acid herbicides, such as [(4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic acid (fluroxypyr).

It is apparent that various modifications may be made in the formulation and application of the compounds of this invention without departing from the inventive concepts herein as defined in the claims.

Table 1
 HERBICIDAL 3-SUBSTITUTED-1-SUBSTITUTED-6-TRIFLUORO-
 METHYL-2,4-(1H,3H)PYRIMIDINEDIONES




Cmpd. No.	R	R ¹	R ²	X	Y	Z
1	t-butyl	-CH ₃	H	O	H	4-Cl
2	t-butyl	-CH ₃	Cl	O	H	4-Cl
3	t-butyl	-NH ₂	H	O	H	4-Cl
4	t-butyl	-CH ₃	H	O	4-Cl	6-F
5	t-butyl	-NH ₂	H	O	4-Cl	6-F
6	-CH ₃	-CH ₃	H	O	4-Cl	6-F
7	Cl	-CH ₃	H	S	H	4-Cl
8	-SCH ₃	-CH ₃	H	S	H	4-Cl
9		-CH ₃	H	S	H	4-Cl

Table 1 (continued)


Cmpd. No.	R	R ¹	R ²	X	Y	Z
10	Cl	-CH ₃	H	S	H	6-F
11	Cl	-CH ₃	Cl	S	H	6-F
12	-CH ₃	-CH ₃	H	S	4-Cl	6-F
13	isopropyl	-CH ₃	H	S	4-Cl	6-F
14	-NHC ₄ H ₉	-CH ₃	H	S	4-Cl	6-F
15		-CH ₃	H	S	4-Cl	6-F
16	-CH ₃	-CH ₃	H	S	H	4-Cl
17	n-propyl	-CH ₃	H	S	H	4-Cl
18	Cl	-CH ₃	H	S	4-OCH ₃	6-Cl
19	isobutyl	-CH ₃	H	O	4-Cl	6-F
20	-CH ₂ OCH ₃	-CH ₃	H	O	4-Cl	6-F
21	-CH ₂ Cl	-CH ₃	H	O	4-Cl	6-F
22	phenyl	-CH ₃	H	O	4-Cl	6-F
23	isopropyl	-CH ₃	H	O	4-Cl	6-F
24	benzyl	-CH ₃	H	O	4-Cl	6-F

Table 1 (continued)






Cmpd. No.	R	R ¹	R ²	X	Y	Z
25	-CH ₃	-CH ₃	H	O	4-Cl	6-Cl
26	hexyl	-CH ₃	H	O	4-Cl	6-F
27	-C(CH ₃) ₂ CH ₂ Cl	-CH ₃	H	O	4-Cl	6-F
28	t-butyl	-CH ₃	H	O	4-Cl	6-Cl
29		-CH ₃	H	O	4-Cl	6-F
30		-CH ₃	H	O	4-Cl	6-F
31		-CH ₃	H	O	4-Cl	6-F
32	t-butyl	-CH ₃	H	O	4-Br	6-F
33	t-butyl	-CH ₃	H	O	4-NO ₂	6-F
34		-CH ₃	H	O	4-Cl	6-F
35		-CH ₃	H	O	4-Cl	6-F

Table 1 (continued)








Cmpd. No.	R	R ¹	R ²	X	Y	Z
36		-CH ₃	H	O	4-Cl	6-F
37		-CH ₃	H	O	4-Cl	6-F
38		-CH ₃	H	O	4-Cl	6-F
39		-CH ₃	H	O	4-Cl	6-F
40		-CH ₃	H	O	4-Cl	6-F
41		-CH ₃	H	O	4-Cl	6-F
42		-CH ₃	H	O	4-Cl	6-F

Table 1 (continued)





Cmpd. No.	R	R ¹	R ²	X	Y	Z
43		-CH ₃	H	O	4-Cl	6-F
44		-CH ₃	H	O	4-Cl	6-F
45	t-butyl	-CH ₃	H	O	H	6-F
46		-CH ₃	H	S	4-Cl	6-F
47	-SC ₃ H ₇	-CH ₃	H	S	4-Cl	6-F
48	-S(O)C ₃ H ₇	-CH ₃	H	S	4-Cl	6-F
49		-CH ₃	H	S	4-Cl	6-F
50	isopropyl	-C-CN	H	S	4-Cl	6-F
51	-SCH ₃	-CH ₃	H	O	4-Cl	6-F
52	-SC ₂ H ₅	-CH ₃	H	O	4-Cl	6-F
53	-SH	-CH ₃	H	O	4-Cl	6-F
54	-SO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
55	-CF ₃	-CH ₃	H	O	4-Cl	6-F

Table 1 (continued)









Cmpd. No.	R	R ¹	R ²	X	Y	Z
56	-CHBr ₂	-CH ₃	H	O	4-Cl	6-F
57	-CH ₂ Br	-CH ₃	H	O	4-Cl	6-F
58		-CH ₃	H	O	4-Cl	6-F
59		-CH ₃	H	O	4-Cl	6-F
60		-CH ₃	H	O	4-Cl	6-F
61		-CH ₃	H	O	4-Cl	6-F
62		-CH ₃	H	O	4-Cl	6-F
63		-CH ₃	H	O	4-Cl	6-F
64		-CH ₃	H	O	4-Cl	6-F
65		-CH ₃	H	O	4-Cl	6-F

Table 1 (continued)

Cmpd. No.	R	R ¹	R ²	X	Y	Z
66	ethyl	-CH ₃	H	O	4-Cl	6-F
67	n-propyl	-CH ₃	H	O	4-Cl	6-F
68	n-butyl	-CH ₃	H	O	4-Cl	6-F
69	sec-butyl	-CH ₃	H	O	4-Cl	6-F
70	-NH ₂	-CH ₃	H	O	4-Cl	6-F
71	-NHSO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
72	-NHSO ₂ C ₆ H ₅	-CH ₃	H	O	4-Cl	6-F
73	-NHSO ₂ NHCH ₃	-CH ₃	H	O	4-Cl	6-F
74	-NHSO ₂ CH ₂ Cl	-CH ₃	H	O	4-Cl	6-F
75	-NHSO ₂ NHC ₆ H ₅	-CH ₃	H	O	4-Cl	6-F
76	-N(CH ₃)SO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
77	-N(C ₂ H ₅)SO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
78	-N(CH ₂ OCH ₃)SO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
79	-N(CH ₂ CH=CH ₂)SO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
80	-N(CH ₂ C≡CH)SO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
81	-N(CH ₃)SO ₂ C ₆ H ₅	-CH ₃	H	O	4-Cl	6-F
82	-NHCH ₂ CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
83	-CO ₂ H	-CH ₃	H	O	4-Cl	6-F
84	-CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F

Table 1 (continued)


Cmpd. No.	R	R ¹	R ²	X	Y	Z
85	-CO ₂ Na	-CH ₃	H	O	4-Cl	6-F
86	-CONH ₂	-CH ₃	H	O	4-Cl	6-F
87	-CONHCH ₃	-CH ₃	H	O	4-Cl	6-F
88	-CON(CH ₃) ₂	-CH ₃	H	O	4-Cl	6-F
89	-CONHSO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
90	-CO ₂ C ₂ H ₅	-CH ₃	H	O	4-Cl	6-F
91		-CH ₃	H	O	4-Cl	6-F
92	-CH ₂ CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
93	-CH ₂ CO ₂ C ₂ H ₅	-CH ₃	H	O	4-Cl	6-F
94	-CH(CH ₃)CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
95	-CH ₂ CH(Cl)CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
96	-CH ₂ CH(CH ₃)CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
97	-CH=C(CH ₃)CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
98	-CH=C(Cl)CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
99	-OCH ₂ CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
100	-OCH(CH ₃)CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
101	-OCH ₂ CH=CH ₂	-CH ₃	H	O	4-Cl	6-F
102	-OCH ₂ C≡CH	-CH ₃	H	O	4-Cl	6-F

Table 1 (continued)








Cmpd. No.	R	R ¹	R ²	X	Y	Z
103	-OH	-CH ₃	H	O	4-Cl	6-F
104	-OCH ₃	-CH ₃	H	O	4-Cl	6-F
105	-OCH(CH ₃) ₂	-CH ₃	H	O	4-Cl	6-F
106	Cl	-CH ₃	H	O	4-Cl	6-F
107	-SCH ₂ CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
108	-SCH ₂ CN	-CH ₃	H	O	4-Cl	6-F
109	-SCH ₂ C≡CH	-CH ₃	H	O	4-Cl	6-F
110	-SCH ₂ C ₆ H ₅	-CH ₃	H	O	4-Cl	6-F
111	-CH ₂ SCH ₂ CO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
112	-CH ₂ SCH ₂ C ₆ H ₅	-CH ₃	H	O	4-Cl	6-F
113	-CH ₂ NHSO ₂ CH ₃	-CH ₃	H	O	4-Cl	6-F
114	-NHPO(OCH ₃) ₂	-CH ₃	H	O	4-Cl	6-F
115	-NHCOCH ₃	-CH ₃	H	O	4-Cl	6-F
116	-N(SO ₂ CH ₃)COCH ₃	-CH ₃	H	O	4-Cl	6-F
117	-CH ₂ OCH ₂ C≡CH	-CH ₃	H	O	4-Cl	6-F
118		-CH ₃	H	O	4-Cl	6-F
119		-CH ₃	H	O	4-Cl	6-F

Table 1 (continued)

Cmpd. No.	R	R ¹	R ²	X	Y	Z
120		-CH ₃	H	O	4-Cl	6-F
121		-CH ₃	H	O	4-Cl	6-F
122		-CH ₃	H	O	4-Cl	6-F
123		-CH ₃	H	O	4-Cl	6-F
124		-CH ₃	H	O	4-Cl	6-F

- 76 -

Table 2
CHARACTERIZING PROPERTIES

<u>Cmpd No</u>	<u>Melting Point (°C)</u>	<u>Cmpd No</u>	<u>Melting Point (°C)</u>
1	146-148	34	oil
2	83-86	35	50-54
3	104-106	36	170-175
4	68-70	37	63-66
5	90-92	38	60-63
6	64-66	39	55-60
7	86-88	40	137-139
8	79-81	41	58-62
9	102-104	42	183-186
10	59-61	43	50-58
11	81-83	44	59-63
12	83-85	45	144-146
13	oil	46	72-74
14	109-111	47	52-54
15	180-181	48	72-75
16	94-96	49	150-153
17	146-148	50	78-80
18	185-187	51	157-158
19	51-53	52	56 °C sublimes
20	54-56	53	86 °C sublimes
21	154-156	55	63-71
22	80-82	56	144-146
23	49-51	58	64-74
24	low melting solid	62	70-75
25	76-78	66	51-59
26	oil	67	43-50
27	65-69	68	47-51
28	64-67	69	50-54
29	50-52	70	solid
30	60-64	93	56-58
31	68-71	105	69-71
32	72-74	124	63-65

- 77 -

Table 3
PREEMERGENCE HERBICIDAL ACTIVITY (% CONTROL)

<u>Compound No.</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	100*	100	100
Wheat	100	98*	100	100
Corn	100	98*	100	100
Velvetleaf	100	100*	100	100
Morningglory	100	100*	100	100
Chickweed	100	100*	100	100
Cocklebur	100	90*	100	100
Blackgrass	95	100*	100	100
Green foxtail	100	100*	100	100
Johnsongrass	100	100*	100	100

<u>Compound No.</u>	<u>5</u>	<u>6</u>	<u>7</u>	<u>8</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	100	60	100
Wheat	95	100	50	80
Corn	100	100	70	90
Velvetleaf	100	100	100	100
Morningglory	100	100	100	95
Chickweed	100	100	100	100
Cocklebur	100	100	100	80
Blackgrass	ND	100	60	90
Green foxtail	100	100	100	100
Johnsongrass	100	100	100	100

<u>Compound No.</u>	<u>9</u>	<u>10</u>	<u>11</u>	<u>12</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	50	90	100
Wheat	85	30	20	100
Corn	95	70	70	100
Velvetleaf	100	100	100	100
Morningglory	100	90	70	100
Chickweed	100	50	0	ND
Cocklebur	50	90	40	100
Blackgrass	80	70	30	ND
Green foxtail	100	100	95	100
Johnsongrass	100	95	100	100

Table 3 (continued)

<u>Compound No.</u>	<u>13</u>	<u>14</u>	<u>15</u>	<u>16</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	100	100	100
Wheat	90	60	80	100
Corn	100	90	95	100
Velvetleaf	100	100	100	100
Morningglory	100	95	100	100
Chickweed	100	100	100	100
Cocklebur	100	50	100	100
Blackgrass	ND	60	95	100
Green foxtail	100	100	100	100
Johnsongrass	100	100	100	100
<u>Compound No.</u>	<u>17</u>	<u>18</u>	<u>19</u>	<u>20</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	20	100	100
Wheat	100	50	80	100
Corn	100	60	90	100
Velvetleaf	100	100	100	100
Morningglory	100	90	100	100
Chickweed	ND	100	100	100
Cocklebur	100	10	100	100
Blackgrass	100	0	100	100
Green foxtail	100	100	100	100
Johnsongrass	100	40	100	100
<u>Compound No.</u>	<u>21</u>	<u>22</u>	<u>23</u>	<u>24</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	50	45	100	95
Wheat	60	30	100	50
Corn	60	50	100	80
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	80	95	100	100
Cocklebur	70	30	100	75
Blackgrass	50	75	100	90
Green foxtail	95	100	100	100
Johnsongrass	90	90	100	100

Table 3 (continued)

<u>Compound No.</u>	<u>25</u>	<u>26</u>	<u>27</u>	<u>28</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	100	100	95
Wheat	90	40	60	55
Corn	95	70	90	90
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	95	100	100	40
Cocklebur	70	100	80	40
Blackgrass	90	60	90	70
Green foxtail	100	100	100	100
Johnsongrass	100	60	100	100

<u>Compound No.</u>	<u>29</u>	<u>30</u>	<u>31</u>	<u>32</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	100	100	100
Wheat	40	40	40	90
Corn	30	20	30	95
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	100	100
Cocklebur	60	60	50	80
Blackgrass	40	60	50	100
Green foxtail	100	100	100	100
Johnsongrass	80	70	60	100

<u>Compound No.</u>	<u>34</u>	<u>35</u>	<u>36</u>	<u>37</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	100	100	100
Wheat	50	60	50	70
Corn	50	60	60	90
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	95	100	100	100
Cocklebur	50	80	90	100
Blackgrass	50	50	60	ND
Green foxtail	100	100	100	100
Johnsongrass	80	75	80	100

Table 3 (continued)

<u>Compound No.</u>	<u>38</u>	<u>39</u>	<u>40</u>	<u>41</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	70	100	100
Wheat	70	50	40	60
Corn	80	40	60	80
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	100	100
Cocklebur	100	100	100	100
Blackgrass	ND	ND	ND	ND
Green foxtail	100	100	100	100
Johnsongrass	95	90	80	95
<u>Compound No.</u>	<u>42</u>	<u>43</u>	<u>44</u>	<u>45</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	70	100	100	100
Wheat	50	65	50	80
Corn	60	50	70	100
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	ND	90
Cocklebur	80	85	100	95
Blackgrass	ND	ND	ND	100
Green foxtail	100	100	100	100
Johnsongrass	80	80	90	100
<u>Compound No.</u>	<u>46</u>	<u>47</u>	<u>48</u>	<u>49</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	100	40	70
Wheat	30	10	60	60
Corn	30	20	80	80
Velvetleaf	100	80	100	100
Morningglory	100	80	100	100
Chickweed	100	0	90	100
Cocklebur	40	80	85	95
Blackgrass	50	ND	ND	ND
Green foxtail	100	100	100	100
Johnsongrass	70	80	75	85

Table 3 (continued)

<u>Compound No.</u>	<u>50</u>	<u>51</u>	<u>52</u>	<u>53</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	70	100	100	100
Wheat	80	90	70	90
Corn	70	100	95	90
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	100	100
Cocklebur	95	100	100	100
Blackgrass	ND	100	95	80
Green foxtail	100	100	100	100
Johnsongrass	80	100	100	100
<u>Compound No.</u>	<u>55</u>	<u>56</u>	<u>58</u>	<u>62</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	20	100	100	100
Wheat	65	80	100	85
Corn	75	90	95	100
Velvetleaf	90	100	100	100
Morningglory	75	100	100	100
Chickweed	90	100	100	100
Cocklebur	50	100	100	100
Blackgrass	50	80	100	95
Green foxtail	95	100	100	100
Johnsongrass	70	90	100	100
<u>Compound No.</u>	<u>66</u>	<u>67</u>	<u>68</u>	<u>69</u>
<u>Rate(kg/ha)</u>	<u>0.1</u>	<u>0.1</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	100	100	100
Wheat	95	90	90	95
Corn	100	95	95	100
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	100	100
Cocklebur	100	100	100	100
Blackgrass	95	100	95	100
Green foxtail	100	100	100	100
Johnsongrass	100	100	95	100

Table 3 (continued)

<u>Compound No.</u>	<u>70</u>	<u>93</u>	<u>124</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>			
Soybean	100	100	100
Wheat	80	100	65
Corn	100	100	80
Velvetleaf	100	100	100
Morningglory	100	100	100
Chickweed	100	100	100
Cocklebur	100	100	100
Blackgrass	95	ND	95
Green foxtail	100	100	100
Johnsongrass	100	100	90

* = Average of two results

ND = no data

Table 4
POSTEMERGENCE HERBICIDAL ACTIVITY (% CONTROL)

<u>Compound No.</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	100*	100	100
Wheat	100	80*	100	100
Corn	100	100*	100	100
Velvetleaf	100	100*	100	100
Morningglory	100	100*	100	100
Chickweed	100	100	100	100
Cocklebur	100	100*	100	100
Blackgrass	100	92*	100	100
Green foxtail	100	100*	100	100
Johnsongrass	100	100*	100	100

<u>Compound No.</u>	<u>5</u>	<u>6</u>	<u>7</u>	<u>8</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	100	80	90
Wheat	100	95	50	85
Corn	100	100	70	85
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	100	100
Cocklebur	100	100	100	100
Blackgrass	ND	ND	60	70
Green foxtail	100	100	100	100
Johnsongrass	100	100	100	100

<u>Compound No.</u>	<u>9</u>	<u>10</u>	<u>11</u>	<u>12</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	90	70	60	100
Wheat	50	20	40	100
Corn	100	90	60	100
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	0	100	ND
Cocklebur	100	100	100	100
Blackgrass	100	0	10	ND
Green foxtail	100	100	100	100
Johnsongrass	100	95	50	100

Table 4 (continued)

<u>Compound No.</u>	<u>13</u>	<u>14</u>	<u>15</u>	<u>16</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	95	95	90	95
Wheat	90	50	70	70
Corn	95	90	95	60
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	100	ND
Cocklebur	100	100	100	90
Blackgrass	ND	50	100	70
Green foxtail	100	100	100	100
Johnsongrass	100	100	100	95
<u>Compound No.</u>	<u>17</u>	<u>18</u>	<u>19</u>	<u>20</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	90	30	90	100
Wheat	100	40	90	100
Corn	100	10	100	100
Velvetleaf	100	100	100	100
Morningglory	90	80	100	100
Chickweed	100	100	100	100
Cocklebur	90	20	ND	100
Blackgrass	95	0	100	100
Green foxtail	100	40	100	100
Johnsongrass	100	50	100	100
<u>Compound No.</u>	<u>21</u>	<u>22</u>	<u>23</u>	<u>24</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	95	90	100	95
Wheat	50	40	100	70
Corn	65	70	100	100
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	90	85	100	100
Cocklebur	80	100	100	100
Blackgrass	50	60	100	90
Green foxtail	100	100	100	100
Johnsongrass	90	80	100	100

Table 4 (continued)

<u>Compound No.</u>	<u>25</u>	<u>26</u>	<u>27</u>	<u>28</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	95	95	100	95
Wheat	70	40	100	65
Corn	80	95	100	80
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	55	100	100	60
Cocklebur	100	100	100	80
Blackgrass	80	70	100	75
Green foxtail	100	100	100	100
Johnsongrass	100	95	100	100

<u>Compound No.</u>	<u>29</u>	<u>30</u>	<u>31</u>	<u>32</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	95	100	95	95
Wheat	50	40	30	95
Corn	100	100	70	100
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	100	100
Cocklebur	100	100	100	100
Blackgrass	80	70	60	100
Green foxtail	100	100	100	100
Johnsongrass	95	90	80	100

<u>Compound No.</u>	<u>34</u>	<u>35</u>	<u>36</u>	<u>37</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	95	95	95	95
Wheat	60	55	60	95
Corn	100	90	90	100
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	100	100
Cocklebur	100	100	100	100
Blackgrass	60	80	70	ND
Green foxtail	100	100	100	100
Johnsongrass	80	80	80	100

Table 4 (continued)

<u>Compound No.</u>	<u>38</u>	<u>39</u>	<u>40</u>	<u>41</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	95	100	100	90
Wheat	60	60	45	80
Corn	100	80	90	100
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	100	100
Cocklebur	100	100	100	80
Blackgrass	ND	ND	ND	ND
Green foxtail	100	100	100	100
Johnsongrass	90	90	80	100
<u>Compound No.</u>	<u>42</u>	<u>43</u>	<u>44</u>	<u>45</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	95	95	95	95
Wheat	50	50	40	70
Corn	90	80	60	80
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	100	70
Cocklebur	100	100	100	70
Blackgrass	ND	ND	ND	70
Green foxtail	100	100	100	100
Johnsongrass	100	95	50	100
<u>Compound No.</u>	<u>46</u>	<u>47</u>	<u>48</u>	<u>49</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	80	95	90	80
Wheat	40	40	50	50
Corn	90	95	80	75
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	ND	80	75
Cocklebur	100	100	100	95
Blackgrass	50	ND	ND	ND
Green foxtail	100	100	100	100
Johnsongrass	70	70	100	90

Table 4 (continued)

<u>Compound No.</u>	<u>50</u>	<u>51</u>	<u>52</u>	<u>53</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	85	95	95	100
Wheat	70	80	70	100
Corn	90	90	95	90
Velvetleaf	100	100	100	100
Morningglory	80	100	100	100
Chickweed	100	100	100	100
Cocklebur	80	100	100	100
Blackgrass	ND	100	100	100
Green foxtail	100	100	100	100
Johnsongrass	100	100	100	100
<u>Compound No.</u>	<u>55</u>	<u>56</u>	<u>58</u>	<u>62</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	70	95	100	100
Wheat	30	50	100	100
Corn	70	70	100	100
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	80	80	100	100
Cocklebur	100	100	100	100
Blackgrass	40	60	100	100
Green foxtail	100	75	100	100
Johnsongrass	75	80	100	100
<u>Compound No.</u>	<u>66</u>	<u>67</u>	<u>68</u>	<u>69</u>
<u>Rate(kg/ha)</u>	<u>0.1</u>	<u>0.1</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>				
Soybean	100	100	95	100
Wheat	100	100	100	100
Corn	100	100	100	100
Velvetleaf	100	100	100	100
Morningglory	100	100	100	100
Chickweed	100	100	100	100
Cocklebur	100	100	100	100
Blackgrass	100	100	100	100
Green foxtail	100	100	100	100
Johnsongrass	100	100	100	100

Table 4 (continued)


<u>Compound No.</u>	<u>70</u>	<u>93</u>	<u>124</u>
<u>Rate(kg/ha)</u>	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>
<u>Species</u>			
Soybean	100	100	95
Wheat	90	100	55
Corn	100	100	100
Velvetleaf	100	100	100
Morningglory	100	100	100
Chickweed	100	100	100
Cocklebur	100	100	100
Blackgrass	90	ND	90
Green foxtail	100	100	100
Johnsongrass	100	100	100

* = Average of two results

ND = no data

Claims

1. A compound of the formula



in which:

R is halogen, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkylarylalkyl, haloalkyl, hydroxy, alkoxy, hydroxyalkyl, haloaryl, haloarylalkyl, alkoxyaryl, sulfhydryl, alkylthio, piperidiny, alkylamino, alkoxyalkyl, phenoxy, amino, alkylsulfonylamino, arylsulfonylamino, carboxy, propionyl, halopropionyl, allyloxy, propargyloxy, acetylamino, alkylthienyl, alkoxyarylalkyl, alkylsulfinyl, alkylsulfonyl, acetoxylalkyl, alkylcarbonyldioxyalkyl, alkylaminosulfonylamino, haloalkylsulfonylamino, arylaminosulfonylamino, N-alkylsulfonyl-N-alkylamino, N-alkylsulfonyl-N-alkoxyalkylamino, N-alkylsulfonyl-N-alkynylamino, N-alkylsulfonyl-N-alkenylamino, N-arylsulfonyl-N-alkylamino, acetoxylalkylamino, acetoxyl, sodium carboxylato, aminocarboxylato, alkylcarbamoyl, alkylsulfonylcarbamoyl, alkoxy carbonyl, acetoxylhaloalkyl, acetoxylalkenyl, acetoxylhaloalkenyl, acetoxylalkoxy, alkenyloxy, alkynyloxy, acetoxylalkylthio, cyanoalkylthio, alkynylthio, arylalkylthio, acetoxylalkylthioalkyl, arylalkylthioalkyl, alkylsulfonylaminoalkyl, alkoxyphosphinyloxyamino, N-acetyl-N-alkylsulfonylamino, alkynyloxyalkyl, alkylcarboxylatophenoxy, halophenoxy, alkylphenoxy, alkoxypropionyloxyphenoxy, or haloalkoxypropionyl;

R¹ is alkyl or amino;

R² is hydrogen or halogen;

X is oxygen or sulfur;

Y is hydrogen, halogen, alkoxy, cyano, or nitro, and;

Z is halogen;

where halogen is bromine, chlorine, fluorine, or iodine, each alkyl moiety has one to six carbon atoms, each aryl moiety is optionally substituted with halogen, alkyl, or alkoxy.

2. A compound of claim 1 in which

R is alkyl, arylalkyl, haloalkyl, hydroxyalkyl, sulfhydryl, alkylthio, piperidiny, amino, alkylamino, or alkoxyalkyl;

R¹ is methyl or amino;

R² is hydrogen or chlorine;

Y is hydrogen, 4-chloro, 4-bromo, or 4-nitro;

Z is 6-chloro when Y is hydrogen or 6-fluoro when Y is other than hydrogen; and

each alkyl moiety has one to four carbon atoms.

3. A compound of claim 2 in which R is alkyl, haloalkyl, hydroxyalkyl, sulfhydryl, alkylthio, alkoxyalkyl, or benzyl, optionally ring-substituted with chlorine or alkoxy; R¹ is methyl; R² is hydrogen; X is oxygen; Y is 4-chloro; and Z is 6-fluoro.

4. A compound of claim 3 in which R is alkyl, dibromomethyl, hydroxyalkyl, sulfhydryl, alkylthio, methoxymethyl, or benzyl ring-substituted with chlorine or methoxy.

5. The compound of claim 4 in which R is t-butyl.

6. The compound of claim 4 in which R is methyl.

7. The compound of claim 4 in which R is methoxymethyl.

8. The compound of claim 4 in which R is 3-methoxybenzyl.

9. The compound of claim 4 in which R is amino.

10. The compound of claim 4 in which R is 1-hydroxy-1-methylethyl.

11. The compound of claim 4 in which R is 2-hydroxy-1,1-dimethylethyl.

12. The compound of claim 4 in which R is dibromomethyl.

13. The compound of claim 4 in which R is sulfhydryl.

14. The compound of claim 4 in which R is methylthio.


15. The compound of claim 4 in which R is ethylthio.

- 91 -

16. A herbicidal composition comprising an herbicidally effective amount of a compound of claim 1 in admixture with an agriculturally acceptable carrier.

17. The method of controlling undesired plant growth which comprises applying to the locus where control is desired a herbicidally effective amount of a composition of claim 16.

18. A compound of the formula



in which:

P is nitro, amino, or azido;

W is hydrogen, alkoxy, or hydroxy;

R¹ is hydrogen, lower alkyl, or amino;

X is oxygen;

Y is hydrogen, hydrogen, halogen, alkoxy, or nitro;

Z is halogen;

and P and W together may be -N-C(O)-O- to complete a benzoxazolinone ring.

19. The compound of claim 18 which is 3-(2-oxo-4-chloro-6-fluoro-2,3-dihydrobenzoxazol-7-yl)-1-methyl-6-trifluoromethyl-2,4-(1H,3H)pyrimidinedione.

INTERNATIONAL SEARCH REPORT

International Application No
PC/US 96/13995

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D413/04 C07D417/04 C07D239/54 A01N43/78 A01N43/76

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 255 047 A (HOFFMANN LA ROCHE) 3 February 1988 see page 18 - page 20; claims 20-22	18
A	see claims	1-17
X	EP 0 408 382 A (NISSAN CHEMICAL IND LTD) 16 January 1991 see page 104 - page 109; claims 1,5-7 see page 118 - page 120	1-19

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "A" document member of the same patent family

Date of the actual completion of the international search

27 November 1996

Date of mailing of the international search report

03.12.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Henry, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC 1/US 96/13995

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0255047	03-02-88	AU-B- 604250	13-12-90
		AU-A- 7637187	11-02-88
		CA-A- 1286662	23-07-91
		US-A- 4859229	22-08-89
		JP-A- 63041466	22-02-88
		ZA-A- 8705466	02-02-88

EP-A-0408382	16-01-91	AT-T- 143017	15-10-96
		AU-B- 627906	03-09-92
		AU-A- 5884990	17-01-91
		CA-A- 2021005	15-01-91
		DE-D- 69028582	24-10-96
		ES-T- 2091799	16-11-96
		US-A- 5084084	28-01-92
		US-A- 5127935	07-07-92
		US-A- 5154755	13-10-92
		JP-A- 3204865	06-09-91

L12 ANSWER 44 OF 74 CAPLUS COPYRIGHT 2005 ACS on STN
 AN 1997:286377 CAPLUS
 DN 126:264105
 TI Herbicidal 3-(substituted benzoxazol-7-yl) and 3-(substituted
 benzothiazol-7-yl)-1-substituted-6-trifluoromethyl-2,4-
 (1H,3H)pyrimidinediones
 IN Crawford, Scott D.; Maravetz, Lester L.; Theodoridis, George
 PA FMC Corp., USA
 SO PCT Int. Appl., 94 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9708170	A1	19970306	WO 1996-US13995	19960830
	W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN				
	AU 9669618	A1	19970319	AU 1996-69618	19960830
	ZA 9610644	A	19970624	ZA 1996-10644	19961218
PRAI	US 1995-3080P	P	19950831		
	WO 1996-US13995	W	19960830		

OS MARPAT 126:264105

AB Herbicidal title compds., compns. containing them, and methods of using them to control undesired plant growth are disclosed, as are novel intermediates used in their preparation The herbicidal compds. are defined as I [R = halo, alk(en/yn)yl, aryl, arylalkyl, alkylarylalkyl, haloalkyl, OH, alkoxy, hydroxyalkyl, haloaryl, haloarylalkyl, alkoxyaryl, SH, alkylthio, piperidinyl, alkylamino, alkoxyalkyl, PhO, amino, alkylsulfonylamino, arylsulfonylamino, CO₂H, etc.; R₁ = alkyl or amino; R₂ = H or halo; X = O or S; Y = H, halo, alkoxy, cyano, or NO₂; and Z = halo; where halo = Br, Cl, F, or iodine, and each alkyl moiety has 1-6 C atoms]. A list of 124 possible specific compds. is given, with phys. and biol. data for over 60 compds. For instance, 3-chloro-4-fluoroaniline reacted with trimethylacetic anhydride to give the corresponding amide, which was lithiated with BuLi and treated with CO₂ to give 2-(tert-butyl)-6-fluorobenzoxazole-7-carboxylic acid. This acid was treated with ClCO₂Et and then NaN₃ to give the acyl azide, which was thermolyzed in refluxing EtOH to give the benzoxazole carbamate derivative II [Y = H]. This was chlorinated by N,N-dichlorourethane in concentrated HCl-AcOH to give II [Y = Cl], which underwent cyclocondensation with CF₃(H₂N)C:CHCO₂Et and then N-methylation with MeI and K₂CO₃ to give title compound III. At 0.3 kg/ha post- or preemergence, III gave nonselective 100% control of all 10 test species, including 3 crops.


IT 188788-12-7P 188788-14-9P 188788-16-1P
 188788-18-3P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(intermediate; preparation of herbicidal benzoxazolyl- and benzothiazolyl-substituted (trifluoromethyl)pyrimidinediones)


RN 188788-12-7 CAPLUS

CN 2,4(1H,3H)-Pyrimidinedione, 3-(3-amino-4-chloro-6-fluoro-2-methoxyphenyl)-6-(trifluoromethyl)- (9CI) (CA INDEX NAME)




RN 188788-14-9 CAPLUS

CN 2,4(1H,3H)-Pyrimidinedione, 3-(3-amino-4-chloro-6-fluoro-2-methoxyphenyl)-1-methyl-6-(trifluoromethyl)- (9CI) (CA INDEX NAME)



RN 188788-16-1 CAPLUS

CN 2,4(1H,3H)-Pyrimidinedione, 3-(3-amino-4-chloro-6-fluoro-2-hydroxyphenyl)-1-methyl-6-(trifluoromethyl)- (9CI) (CA INDEX NAME)



RN 188788-18-3 CAPLUS

CN Propanedioic acid, 3-chloro-6-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-2-[(3-ethoxy-1,3-dioxopropyl)amino]-5-hydroxyphenyl ethyl ester (9CI) (CA INDEX NAME)

